

Markov Processes – Problem Sheet 5.

Tutorials by Nikolay Barashkov <s6nibara@uni-bonn.de>, Robert Crowell <crowellr@googlemail.com>. Solutions will be collected Tuesday November 28st during the lecture. At most in groups of 2.

The goal of this problem sheet is to apply Lyapunov functions to study diffusion processes on \mathbb{R}^d and Markov chains corresponding to Euler approximations of the corresponding stochastic differential equations.

Let $b: \mathbb{R}^d \rightarrow \mathbb{R}^d$ and $\sigma: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d}$ be locally Lipschitz continuous functions. We consider a diffusion process (X_t, \mathbb{P}_x) with possibly finite life-time ζ solving a stochastic differential equation

$$dX_t = \sigma(X_t)dB_t + b(X_t)dt \quad \text{for } t < \zeta, X_0 = x,$$

where (B_t) is a d -dimensional Brownian motion. Let $a(x) = \sigma(x)\sigma(x)^T$. For the exercises below, it will only be important to know that (X_t, \mathbb{P}_x) solves the local martingale problem for the generator

$$\mathcal{L}f(x) = \frac{1}{2} \sum_{i,j} a_{i,j}(x) \nabla^i \nabla^j f(x) + \sum_i b_i(x) \nabla^i f(x),$$

in the sense that for any $f \in C^{1,2}(\mathbb{R}_+ \times \mathbb{R}^d)$,

$$M_t^f = f(X_t) - \int_0^t \mathcal{L}f(s, X_s) ds$$

is a local martingale wrt. \mathbb{P}_x . More precisely, let $T_k = \inf\{t \geq 0: |X_t| \geq k\}$, then $\zeta = \sup_k T_k$ and for any k the stopped process $M_{t \wedge T_k}^f$ is a martingale under \mathbb{P}_x .

For a fixed time step $h > 0$, the *Euler-Maruyama approximation* of the diffusion process above is the time-homogeneous Markov chain (X_n^h, \mathbb{P}_x) with transition step

$$x \mapsto x + b(x)h + \sqrt{h}\sigma(x)Z, \quad Z \sim \mathcal{N}(0, \mathbb{I}_d).$$

We denote the corresponding transition kernel and generator by π_h and \mathcal{L}_h , respectively.

Exercise 1. (EXPLOSIONS)

- a) Prove that the diffusion process is non-explosive if

$$\text{Tr } a(x) / 2 + x \cdot b(x) = O(|x|^2), \quad \text{as } |x| \rightarrow \infty.$$

- b) Show that for $\varepsilon > 0$, the condition $O(|x|^2)$ cannot be replaced by $O(|x|^{2+\varepsilon})$.
- c) Implement the Euler scheme on a computer, e.g. for $d=1$. Do some experiments. Can you observe a different behavior in cases where the condition above is satisfied or violated, respectively?
- d) As discrete time Markov chains, the Euler approximations always have infinite lifetime. Which properties can you prove for the Euler approximations under similar conditions as in a)?

Exercise 2. (STATIONARY DISTRIBUTIONS I) Suppose that $\zeta = \infty$ almost surely, and that there exists $V \in C^{1,2}(\mathbb{R}_+ \times \mathbb{R}^d)$, $\varepsilon, c \in \mathbb{R}_+$ and a ball $B \subset \mathbb{R}^d$ such that $V \geq 0$ and

$$\frac{\partial V}{\partial t} + \mathcal{L}V \leq -\varepsilon + c\mathbb{I}_B, \quad \text{on } \mathbb{R}_+ \times \mathbb{R}^d. \quad (1)$$

a) Prove that

$$\mathbb{E} \left[\frac{1}{t} \int_0^t \mathbb{I}_B(X_s) ds \right] \geq \frac{\varepsilon}{c} - \frac{V(0, x_0)}{ct}.$$

b) It can be shown that (1) implies that (X_t, \mathbb{P}_x) is a time-homogeneous Markov process with Feller transition semigroup $(p_t)_{t \geq 0}$. Conclude that there exists an invariant probability measure μ . *Hint: Try to carry over the proof in discrete time to the continuous time case.*

Exercise 3. (STATIONARY DISTRIBUTIONS II) Suppose that the conditions in Exercise 2 hold, and let μ be an invariant probability measure.

a) Show that $\int \mathcal{L}f d\mu = 0$ for any $f \in C_0^\infty(\mathbb{R}^d)$.

b) Use this to compute μ explicitly in the case $d=1$. Here, assume that b and σ are twice continuously differentiable, and $\sigma(x) > 0$ for all x . You may also assume without proof that μ has a twice continuously differentiable density $\rho(x)$ which is strictly positive.

Exercise 4. (STATIONARY DISTRIBUTIONS III)

a) Show that an invariant probability measure for the diffusion process on \mathbb{R}^d exists if

$$\limsup_{|x| \rightarrow \infty} \left(\frac{1}{2} \text{Tr } a(x) + x \cdot b(x) \right) < 0. \quad (2)$$

b) Give conditions ensuring the existence of an invariant probability measure for the Euler approximations. Why is not enough to assume (2)?

c) Study different cases experimentally using an implementation of the Euler scheme. Can you see the difference between cases where an invariant probability measure exists or does not exist?

Die Fachschaft Mathematik feiert am 23.11. ihre Matheparty in der N8schicht. Der VVK findet am Mo. 20.11., Di. 21.11. und Mi 22.11. in der Mensa Poppelsdorf statt.

Alle weitere Infos auch auf fsmath.uni-bonn.de

The student council of mathematics will organize the math party on 23/11 in N8schicht. The presale will be held on Mon 20/11, Tue 21/11 and Wed 22/11 in the mensa Poppelsdorf.

Further information can be found at fsmath.uni-bonn.de