

## Stochastic Analysis – Problem Sheet 7.

Tutorial classes: Mon July 3rd 16–18 in SemR 1.007. Claudio Bellani <claudio.bellani01@gmail.com>.  
Solutions will be collected Tuesday June 27th during the lecture. At most in groups of 3.

Let  $(\Omega := C(\mathbb{R}_{\geq 0}; \mathbb{R}), \mathcal{F}, \mathcal{F}_\bullet, \mathbb{P})$  the one dimensional Wiener space and  $X$  the canonical process.

**Exercise 1.** Find a predictable process  $F$  such that

$$\Phi = \mathbb{E}[\Phi] + \int_0^\infty F_s dX_s$$

when  $\Phi \in L^2(\Omega, \mathcal{F}_T, \mathbb{P})$  is each of the following r.v. (with  $T > 0$  fixed)

$$X_T^2, \quad e^{X_T}, \quad \int_0^T X_t dt, \quad X_T^3, \quad \sin(X_T).$$

**Exercise 2.** We want to prove that the linear span of r.v. of the form

$$E(h) = \cos\left(\int h_s dX_s\right) \exp\left(\frac{1}{2} \int h_s^2 ds\right), \quad F(h) = \sin\left(\int h_s dX_s\right) \exp\left(\frac{1}{2} \int h_s^2 ds\right), \quad h \in L^2(\mathbb{R}_{\geq 0}),$$

is dense in  $L^2(\Omega, \mathcal{F}, \mathbb{P})$  ( $h$  is a deterministic function and the integrals are over  $\mathbb{R}_{\geq 0}$ ).

a) Show that if  $G \in L^2(\Omega, \mathcal{F}, \mathbb{P})$  is orthogonal to all  $\{E(h), F(h): h \in L^2(\mathbb{R})\}$ , then in particular

$$\mathbb{E}[G \exp(i\lambda_1 B_{t_1} + \dots + i\lambda_n B_{t_n})] = 0$$

for all  $\lambda_1, \dots, \lambda_n \in \mathbb{R}$  and  $t_1, \dots, t_n \geq 0$ .

b) Deduce from this that  $G$  is orthogonal to all functions of the form  $\phi(B_{t_1}, \dots, B_{t_n})$  with  $\phi \in C_0^\infty$ .  
[Hint: use Fourier transform]

c) Conclude.

**Exercise 3.** Use the class of functions introduced in Exercise 2 to prove the Brownian martingale representation theorem.

a) Determine the martingale representation for functions  $\Phi$  of the form

$$\Phi = \sum_i (a_i E(h_i) + b_i F(h_i))$$

where  $a_i, b_i \in \mathbb{R}$ ,  $h_i \in L^2(\mathbb{R}_{\geq 0})$  and the sum is finite.

b) Use the density of such functions to approximate an arbitrary element  $\Phi \in L^2$  and conclude.