

## V4F1 Stochastic Analysis – Problem Sheet 6

Version 1, 2020.06.1. Tutorial classes: Mon June 8th 16–18 (Zoom) Min Liu | Wed June 10th 16–18 (Zoom) Daria Frolova.

Solutions in groups of 2 (at most). To be handled in L<sup>A</sup>T<sub>E</sub>X or T<sub>E</sub>X<sub>M</sub>A<sub>C</sub>S format via eCampus not later than **8pm Friday June 5th**. Use this sheet for your solutions and write them under the corresponding exercise. Fill out your names below.

Names: XXXXXXXXXXXXX/YYYYYYYYYYYYYYYY

**Exercise 1 (Pts 2+2+2+2) (Brownian motion writes your name)** Prove that a Brownian motion in  $\mathbb{R}^2$  will write your name (in cursive script, without dotted 'i's or crossed 't's). Let  $B$  be a two dimensional Brownian motion on  $[0, 1]$  and observe that  $X_t^{(a,b)} = (b-a)^{1/2}(B_{a+(b-a)t} - B_a)$  for  $t \in [0, 1]$  has the same law as  $B$ . Let  $g : [0, 1] \rightarrow \mathbb{R}^2$  a smooth parametrization of your name. Let us agree that the Brownian motion  $X^{(a,b)}$  spells your name (to precision  $\varepsilon > 0$ ) if

$$\sup_{t \in (0,1)} |X_t^{(a,b)} - g(t)| \leq \varepsilon. \quad (1)$$

- a) For  $k \in \mathbb{N}$  let  $A_k$  be the event that (1) holds for  $a = 2^{-k-1}$  and  $b = 2^{-k}$ . Check that the events  $(A_k)_{k \in \mathbb{N}}$  are independent and  $\mathbb{P}(A_k) = \mathbb{P}(A_0)$  for all  $k \geq 0$ . Conclude that if  $\mathbb{P}(A_0) > 0$  then infinitely many of the  $A_k$ s will occur almost surely.
- b) Show that

$$\mathbb{P}[\sup_{t \in (0,1)} |B_t| \leq \varepsilon] > 0. \quad (2)$$

- c) Using (2) and Girsanov's transform to show that  $\mathbb{P}(A_0) > 0$  (Hint: construct a measure  $\mathbb{Q}$  so that  $B_t - g(t)$  is a Brownian motion)
- d) Prove that a similar result holds for  $g$  only continuous.

**Exercise 2 (Pts 3)** Let  $(X, \mathbb{P})$  be a solution of the martingale problem with drift  $b$  and diffusion  $\sigma$ . Generalise appropriately the Girsanov transform to construct a measure  $\mathbb{Q}$  under which the process  $X$  solves a martingale problem with a different drift. For simplicity, assume that all the necessary integrability conditions are satisfied. (What takes the place of the Brownian motion?)

**Exercise 3 (Pts 3+3+3)** Given smooth, bounded functions  $A : \mathbb{R}^d \rightarrow \mathbb{R}^d$ ,  $V : \mathbb{R}^d \rightarrow \mathbb{R}$ . Consider the operator  $H(A)$  on  $L^2(\mathbb{R}^d)$  given by

$$H(A) = -\frac{1}{2}|\nabla - iA(x)|^2 + V(x)$$

We will assume that this operator is self-adjoint (with suitable domain), bounded from below and with discrete spectrum. We will denote  $E_0(A)$  its smaller eigenvalue which we will assume simple (i.e. of multiplicity one). Let  $\psi$  the complex valued solution to

$$\partial_t \psi(t, x) = -H(A)\psi(t, x), \quad \psi(0, x) = \psi_0(x),$$

which we will assume to exist, to be once differentiable in  $t$  and twice in  $x$  and be bounded with bounded derivatives.

- a) Find a suitable functions  $B, C : \mathbb{R}^d \rightarrow \mathbb{C}$  with which we can give the following Feynman–Kac representation for  $\psi$ :

$$\psi(t, x) = \mathbb{E}_x \left\{ \psi_0(X_t) \exp \left[ \int_0^t B(X_s) dX_s + \int_0^t C(X_s) ds \right] \right\}$$

where under  $\mathbb{E}_x$  the process  $X$  is a  $d$ -dimensional Brownian motion starting at  $x \in \mathbb{R}^d$ .

- b) Prove that the lowest eigenvector of  $H_A$  is strictly positive everywhere.
- c) Use the above representation to prove the *diamagnetic inequality*

$$E_0(A) \geq E_0(0).$$

[Hint: take  $\psi_0(x) = 1$  and argue that  $\psi(t, x) \simeq ce^{-E_0 t} \varphi(x) + o_t(1)$  where  $H\varphi = E_0(A)\varphi$  and conclude]