

V4F1 Stochastic Analysis – Problem Sheet 7

Version 2, 2020.06.10. Tutorial classes: Mon June 15th 16–18 (Zoom) Min Liu | Wed June 17th 16–18 (Zoom) Daria Frolova.
 Solutions in groups of 2 (at most). To be handled in L^AT_EX or T_EX_{MACS} format via eCampus not later than **8pm Friday June 12th**. Use this sheet for your solutions and write them under the corresponding exercise. Fill out your names below.

Names: **XXXXXXXXXXXXX/YYYYYYYYYYYYYYY**

Exercise 1 (Pts 3+3+3+4) Let X a solution of the SDE in \mathbb{R}^n

$$dX_t = b(X_t)dt + dB_t, \tag{1}$$

with a vectorfield $b : \mathbb{R}^n \rightarrow \mathbb{R}^n$ measurable and with linear growth.

a) Prove that for all $T > 0$, almost surely

$$A(T) = \int_0^T |b(X_s)|^2 ds < \infty,$$

and therefore the process is unique in law.

b) Find a (deterministic) increasing function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that, almost surely

$$\sup_{T \geq 0} \frac{A(T)}{f(T)} < \infty.$$

[Hint: find a constant C such that $\sup_{T \geq 0} \frac{A(T)}{f(T)} \leq \sum_{n \geq 0} \frac{CA(n)}{f(n)} < \infty$ a.s.]

c) Use Girsanov's transform to prove that the process is Markov when b is a bounded vectorfield.

d) (Bonus) Try to extend the proof of the Markov property for b of linear growth.

Exercise 2 (Pts 5) Let $C^n = C(\mathbb{R}_+, \mathbb{R}^n)$ with the Borel σ -field and \mathbb{W}_x the law of the Brownian motion starting at x . Let X the unique solution of the SDE (1) with $b = -\nabla V$ and V a positive C^2 function such that

$$|\nabla V(x)|^2 - \Delta V(x) \geq -L \quad x \in \mathbb{R}^n.$$

Use the path-integral formula

$$\mathbb{E}_x(f(X_T)) = \int_{C^n} f(\omega_T) \exp\left(V(\omega_0) - V(\omega_T) - \frac{1}{2} \int_0^T (|\nabla V(\omega_s)|^2 - \Delta V(\omega_s)) ds\right) \mathbb{W}_x(d\omega)$$

to show that for any two bounded functions f, g and under appropriate conditions on V :

$$\int (P_T f)(x) g(x) e^{-2V(x)} dx = \int f(x) (P_T g)(x) e^{-2V(x)} dx$$

which shows that P_T is symmetric wrt. the measure $e^{-2V(x)} dx$ and taking $g = 1$ show that $e^{-2V(x)} dx$ properly normalized is an invariant measure for the SDE

$$dX_t = -\nabla V(X_t)dt + dB_t,$$

meaning that if X_0 is taken with probability distribution $\propto e^{-2V(x)} dx$ then

$$\mathbb{E}[f(X_0)] = \mathbb{E}[f(X_T)],$$

for all $T \geq 0$.

[Hint: let $\mathbb{W}_{x,y}$ the conditional law of the Brownian motion ω to have $\omega_T = y$, i.e. the Brownian bridge. Prove that the under $\mathbb{W}_{x,y}$ the process $\tilde{\omega}_t = \omega_{T-t}$ has law $\mathbb{W}_{y,x}$ and use the path integral]

Exercise 3 (Pts 3+3) Prove a Fubini theorem for stochastic integrals. Let (Λ, \mathcal{A}) be a measure space and $(\Omega, \mathcal{F}, \mathcal{F}_\bullet, \mathbb{P})$ a filtered probability space.

- a) Let $(X_n)_n$ a sequence of functions $X_n : \Omega \times \Lambda \rightarrow \mathbb{R}$ which are $\mathcal{F} \otimes \mathcal{A}$ measurable (product σ -field) and such that $(X_n(\cdot, \lambda))_n$ converges in probability for any fixed $\lambda \in \Lambda$. Prove that there exists an $\mathcal{F} \otimes \mathcal{A}$ measurable function $X : \Omega \times \Lambda \rightarrow \mathbb{R}$ for which $X_n(\cdot, \lambda) \xrightarrow{\mathbb{P}} X(\cdot, \lambda)$ for any $\lambda \in \Lambda$. [Hint: here the difficulty is the measurability of the limit X , consider the sequence $n_k(\lambda)$ defined by $n_0(\lambda) = 1$ and

$$n_{k+1}(\lambda) = \inf\{m > n_k(\lambda) : \sup_{p, q \geq m} \mathbb{P}[|X_p(\cdot, \lambda) - X_q(\cdot, \lambda)| > 2^{-k}] \leq 2^{-k}\}$$

and prove that $\lim_k X_{n_k(\lambda)}(\cdot, \lambda)$ exists a.s. and conclude]

- b) Let $H : \Lambda \times \mathbb{R}_{\geq 0} \times \Omega \rightarrow R$ be a bounded function which is measurable w.r.t. $\mathcal{A} \otimes \mathcal{P}$ where \mathcal{P} is the predictable σ -field on $\mathbb{R}_{\geq 0} \times \Omega$. Let M be a continuous martingale on $(\Omega, \mathcal{F}, \mathcal{F}_\bullet, \mathbb{P})$. Prove that there exists a function $J : \Lambda \times \Omega \rightarrow \mathbb{R}$ measurable for $\mathcal{A} \otimes \mathcal{F}_T$ which is a version of the stochastic process $\lambda \mapsto J(\lambda) := \int_0^T H(\lambda, s) dM_s$ and for which it holds

$$\int_{\Lambda} J(\lambda) m(d\lambda) = \int_0^T \left[\int_{\Lambda} H(\lambda, s, \cdot) m(d\lambda) \right] dM_s, \quad a.s.$$

for any bounded measure m on (Λ, \mathcal{A}) . Hint: prove that

$$\mathbb{E} \left[\left(\int_0^T \left[\int_{\Lambda} H(\lambda, s, \cdot) m(d\lambda) \right] dM_s - \int_{\Lambda} J(\lambda) m(d\lambda) \right)^2 \right] = 0.$$

[Taken from Revuz-Yor, Chap. 4]
