


***Stochastic analysis
of quantum fields***





Quantum mechanics

+

Special relativity

=

Quantum field theory


VOLUME I FOUNDATIONS

THE
QUANTUM
THEORY OF
FIELDS

STEVEN WEINBERG

$$(\partial_t^2 - \Delta)\Phi(z) = F(\Phi(z)), \quad z \in \mathbb{R}^{3+1}$$

$$[\Phi(t, x), \dot{\Phi}(t, y)] = \delta(x - y)$$



Φ is an operator valued distribution on an Hilbert space:

$\Phi(f)$ is an unbounded self-adjoint operator for all test functions $f: \mathbb{R}^{3+1} \rightarrow \mathbb{R}$

Does a consistent theory of quantum fields exist?

▷ Axiomatic approach ('60): Wightman, Haag-Kastler.

- i. Hilbert space \mathcal{H} ,
- ii. Positive energy representation $(U(\Lambda, a))_{(\Lambda, a)}$ of Poincaré group $G = \{(\Lambda, a)\}$
- iii. Fields $\varphi(x)$ transforming as $U(\Lambda, a)^* \varphi(x) U(\Lambda, a) = \varphi(\Lambda x + a)$, $x \in \mathbb{R}^{3+1}$
- iv. Vacuum vector $\Omega \in \mathcal{H}$: $U(\Lambda, a)\Omega = \Omega$

▷ Examples?

Progress in \mathbb{R}^{1+1} (1965-1976): Nelson, Glimm, Jaffe, Segal...

▷ Schwinger

$$t \rightarrow it \quad \square \rightarrow \Delta$$

- ▷ Symanzik – functional integral representation of Euclidean correlation function
- ▷ Nelson – reconstruction of the Hilbert space from Euclidean data and spatial Markov property
- ▷ Schwinger functions

$$S_n(f_1 \otimes \cdots \otimes f_n) := \int_{\mathcal{F}'(\mathbb{R}^{d+1})} \varphi(f_1) \cdots \varphi(f_n) \nu(d\varphi).$$

Distribution property – $\beta > 0$,

$$|S_n(f_1 \otimes \dots \otimes f_n)| \leq (n!)^\beta \prod_{i=1}^n \|f_i\|_s. \quad \forall n \geq 0, f_1, \dots, f_n \in \mathcal{S}(\mathbb{R}_+^3).$$

Euclidean invariance – $(a, R).f_n(x) = f_n(a + Rx)$, $(a, R) \in \mathbb{R}^3 \times O(3)$


$$S_n((a, R).f_1 \otimes \dots \otimes (a, R).f_n) = S_n(f_1 \otimes \dots \otimes f_n),$$

Reflection positivity – $(f_n \in \mathcal{S}_{\mathbb{C}}(\mathbb{R}_+^{3n}))_{n \in \mathbb{N}_0}$ (with finitely many nonzero elements)

$$\sum_{n, m \in \mathbb{N}_0} S_{n+m}(\overline{\theta} f_n \otimes f_m) \geq 0,$$

Symmetry – $S_n(f_1 \otimes \dots \otimes f_n) = S_n(f_{\pi(1)} \otimes \dots \otimes f_{\pi(n)})$

- ▷ ('70-'80) Glimm, Jaffe. Nelson. Segal. Guerra, Rosen, Simon, and many others...
- ▷ Construction of theories in $2+1$ dimensions
- ▷ $(\Phi_{\frac{4}{3}}^4)_\Lambda$ Glimm ('69). Glimm, Jaffe. Feldman ('74), Y.M.Park ('75)
- ▷ $(\Phi_{\frac{4}{3}}^4)_{\mathbb{R}^3}$ Feldman, Osterwalder ('76). Magnen, Sénéor ('76). Seiler, Simon ('76)
- ▷ *Other constructions of $\Phi_{\frac{4}{3}}^4$.* Benfatto, Cassandro, Gallavotti, Nicolò, Olivieri, Presutti, Scacciatelli ('80) Brydges, Fröhlich, Sokal ('83) Battle, Federbush ('83) Williamson ('87) Balaban ('83) Gawedzki, Kupiainen ('85) Watson ('89) Brydges, Dimock, Hurd ('95)



Ito and Dœblin want to study diffusion processes via their *sample paths*

Measures

Samples

$$(\mu_t)_t \subseteq \Pi(S)$$

$$\mu_t(dy) = \int P_{t-s}(x, dy) \mu_s(dx)$$

$$X: \Omega \rightarrow C(\mathbb{R}_+, S)$$

$$dX_t = b(X_t)dt + dB_t$$

Advantages

- lower dimensional problem
- more tools (e.g. fixpoint theorems)
- more intuition
- **canonical** reference object $(B_t)_t$

Copyrighted Material

DANIEL REVUZ
MARC YOR


Volume 293

Grundlehren
der mathematischen
Wissenschaften

A Series of
Comprehensive Studies
in Mathematics

CONTINUOUS
MARTINGALES
AND
BROWNIAN MOTION

THIRD EDITION

 Springer

Copyrighted Material

ENCYCLOPEDIA OF MATHEMATICS AND ITS APPLICATIONS 45

STOCHASTIC EQUATIONS IN INFINITE DIMENSIONS

GIUSEPPE DA PRATO
& JERZY ZABCZYK

Relation between a stochastic differential equation and a probability measure

(broadly speaking)

- ▷ Nelson and Parisi–Wu ('84) advocated the *constructive* use of stochastic partial differential equations (SPDEs) to realize a given Gibbs measure for the use of Euclidean quantum field theory (in particular gauge theories)
- ▷ Theoretical version of MCMC methods

$\Lambda = \text{finite set}, \mathbb{T}^d, \mathbb{R}^d$

equation $\partial_t \phi(t) = -\frac{\delta V(\phi(t))}{\delta \phi} + \sqrt{2} \xi(t), \quad \phi: \mathbb{R}_+ \times \Lambda \rightarrow \mathbb{R}$

measure $\phi(t) \sim \nu(d\varphi) = \frac{e^{-V(\varphi)}}{Z} d\varphi \in \text{Prob}(\Lambda \rightarrow \mathbb{R})$

- ▷ The measure ν is described via *white noise*
- ▷ Markov process, invariant measures, ergodicity

$$V(\varphi) = \int \frac{1}{2} |\nabla \varphi|^2 + \frac{m^2 - \infty}{2} \varphi^2 + \frac{\lambda}{4} \varphi^4.$$

$$\partial_t \varphi = \Delta \varphi - \lambda(\varphi^3 - \infty \varphi) - m^2 \varphi + \sqrt{2} \xi \quad \mathbb{R}^3 \times \mathbb{R}_+$$


($d=2$) Jona-Lasinio, P.K.Mitter ('85) Borkar, Chari, S.K.Mitter ('88) Albeverio, Röckner ('91) Da Prato, Debussche ('03) Mourrat, Weber ('17) Tsatsoulis, Weber ('16) Röckner, R.Zhu, X.Zhu ('17)

▷ $d=3$ is more singular: regularity structures (Hairer), paracontrolled distributions (G. Imkeller, Perkowski)

Hairer ('14) Kupiainen ('16) Catellier, Chouk ('17) Mourrat, Weber ('17) Hairer, Mattingly ('18) R.Zhu, X.Zhu ('18) Albeverio, Kusuoka ('18) G. Hofmanová ('18) Moinat, Weber ('18)

Reflection positivity + Euclidean invariance \Rightarrow singularities, infinite volume limit

G. Hofmanová ('18) – construction of $\Phi_{\frac{4}{3}}$ on \mathbb{R}^3 via stochastic quantisation and verification of (most of) the axioms.



- ▷ Much like Ito's approach to diffusions
- ▷ Markovianity does not play any role

equation

$$\left\{ \begin{array}{l} \partial_t \phi(t) = -\frac{\delta H(\phi(t), \dot{\phi}(t))}{\delta \dot{\phi}} \\ \partial_t \dot{\phi}(t) = \underbrace{-\frac{\delta H(\phi(t), \dot{\phi}(t))}{\delta \phi}}_{\text{Hamiltonian dynamics}} \underbrace{-\gamma \dot{\phi}(t) + \sqrt{2} \xi(t)}_{\text{linear Langevin dynamics}}, \end{array} \right. \quad \phi, \dot{\phi}: \mathbb{R} \times \Lambda \rightarrow \mathbb{R}$$

$$H(\varphi, \dot{\varphi}) := V(\varphi) + \frac{\gamma}{2} \dot{\varphi}^2$$

measure

$$(\phi(t), \dot{\phi}(t)) \sim \nu(d\varphi d\dot{\varphi}) = \frac{e^{-H(\varphi, \dot{\varphi})}}{Z} d\varphi d\dot{\varphi} \in \text{Prob}(\Lambda \rightarrow \mathbb{R}^2)$$

▷ Introduced by Ryang, Saito and Shigemoto ('85).

For Φ_d^4 , $d=1,2,3$

$$V(\varphi) = \int \frac{1}{2} |\nabla \varphi|^2 + \frac{m^2 - \infty}{2} \varphi^2 + \frac{\lambda}{4} \varphi^4,$$

$$\partial_t^2 \phi = \Delta \phi + (m^2 - \infty) \phi + \lambda \phi^3 - \gamma \partial_t \phi + \sqrt{2} \xi,$$

Problem: no Schauder estimates, scaling arguments less clear.

Conjecture: *same* renormalization constants of the static measure!

▷ $d=1$. Tolomeo ('18) unique ergodicity.

▷ $d=2$. G, Koch, Oh ('18) local well-posedness (any polynomial), G, Koch, Oh, Tolomeo (in preparation) global well-posedness.

▷ $d=3$. G, Koch, Oh ('18) only quadratic nonlinearity.

equation $\Delta_z \phi(z) = -\frac{\delta V(\phi(z))}{\delta \phi} + \xi(z), \quad \phi: \mathbb{R}^2 \times \Lambda \rightarrow \mathbb{R}$

measure $\phi(z) \sim \nu(d\varphi) = \frac{e^{-4\pi V(\varphi)}}{Z} d\varphi \in \text{Prob}(\Lambda \rightarrow \mathbb{R})$

Discovered perturbatively by Imry, Ma ('75), Young ('77). Non-perturbative “proof” by Parisi and Sourlas ('79-'82) using *supersymmetry*

$$(\text{SPDE})_{d+2} \xrightarrow{\text{“Girsanov”}} (\text{SUSY EQFT})_{d+2} \xrightarrow{\text{dimensional reduction}} (\text{measure})_d$$

$$V(\varphi) = \frac{1}{2} m^2 \varphi^2$$

$$\Delta_z \varphi(z) = -m^2 \varphi(z) + \xi(z), \quad z \in \mathbb{R}^2$$

$$\varphi(z) = \int_{\mathbb{R}^d} \frac{e^{ik \cdot z}}{|k|^2 + m^2} \frac{\eta(dk)}{2\pi}$$

$$\mathbb{E}[\varphi(0)^2] = \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} \frac{dk}{(|k|^2 + m^2)^2} = \frac{1}{(2\pi)^2 m^2} \int_{\mathbb{R}^2} \frac{dk}{(|k|^2 + 1)^2} = \frac{1}{4\pi m^2} \int_0^\infty \frac{d\rho^2}{(\rho^2 + 1)^2} = \frac{1}{4\pi m^2}$$

$$\varphi(0) \sim e^{-4\pi \frac{m^2}{2} \phi^2} d\phi \sim e^{-4\pi V(\phi)} d\phi$$

- ▷ Rigorous proof of dimensional reduction by Klein, Landau and Perez ('84)
- ▷ Recently complete proof by Albeverio, G. and De Vecchi ('18). First for Λ finite dimensional + technical conditions. Then extended to (some) renormalized EQFT.

Stochastic quantisation of Liouville action up to the critical value of $\sigma^2 < 8\pi$ in $\Lambda = \mathbb{T}^2$

$$V(\varphi) = \int_{\mathbb{T}^2} \frac{1}{2} |\nabla \varphi|^2 + \alpha e^{\sigma\varphi - \sigma^2\infty}$$

An alternative approach to induce an equation for $\Phi_{\frac{4}{3}}$ in \mathbb{T}^3 [Barashkov, G. ('18)]

$$e^{-\mathcal{W}(f)} = \int_{\mathcal{P}'(\mathbb{T}^3)} e^{\langle f, \phi \rangle - \lambda \int \phi^4} \underbrace{e^{-\frac{1}{2} \int |\nabla \phi|^2 + a \phi^2}}_{\text{Gaussian free field}} d\phi = \lim_{T \rightarrow \infty} \mathbb{E}[e^{\langle f, W_T \rangle - \lambda \int (W_T^4 - a_T W_T^2)}]$$

▷ W_T is a regularisation of the Gaussian free field at scale T :

$$\mathbb{E}[W_T(f)W_S(f)] = (T \wedge S) \langle f, \rho((-\Delta)^{1/2}/S)(1-\Delta)^{-1}g \rangle$$

$$W_t = \int_0^t J_t dX_t$$

with $(X_t)_t$ cylindrical Wiener process in $L^2(\mathbb{T}^3)$.

Take $V_T(\phi) = \lambda \int (\phi^4 - a_T \phi^2 - b_T)$

$$\mathcal{W}_T(f) = -\log \mathbb{E}[e^{\langle f, W_T \rangle - V_T(W_T)}]$$

$$= \inf_u \mathbb{E} \left[\langle f, W_T + I_T(u) \rangle - V_T(W_T + I_T(u)) + \frac{1}{2} \int_0^\infty \|u_s^2\|_{L^2(\mathbb{T}^3)} ds \right]$$

where

$$I_T(u) = \int_0^T J_t u_t dt$$

and the infimum is over processes u adapted to the Wiener filtration.

$\exists a_T, b_T$ with $a_T, b_T \rightarrow \infty$ as $T \rightarrow \infty$, such that

$$\begin{aligned} \mathcal{W}_T(f) &= \mathbb{E} \left[-\langle f, W_T + I_T(u) \rangle + \lambda V_T(W_T + I_T(u)) + \frac{1}{2} \|u\|_{\mathcal{H}}^2 \right] \\ &= \mathbb{E} \left[-f(W_T + I_T(u)) + \Phi_T(W, u) + \lambda \int (I_T(u))^4 + \frac{1}{2} \|I^T(u)\|_{\mathcal{H}}^2 \right] \end{aligned}$$

$$I_t^T(u) := u_t + \lambda \mathbb{1}_{t \leq T} J_t W_t^3 + \lambda \mathbb{1}_{t \leq T} J_t (W_t^2 \triangleright I_t^b(u)), \quad \mathcal{H} = L^2(\mathbb{R}_+ \times \mathbb{T}^3),$$

$$|\mathbb{E} \Phi_T(W, u)| \leq \mathbb{E} Q(W) + \frac{1}{4} (\lambda \|I_T(u)\|_{L^4}^4 + \|I^T(u)\|_{\mathcal{H}}^2)$$

Barashkov, G. ('18)

Theorem

$$\begin{aligned} \mathcal{W}_{\infty}(f) &:= \liminf_{T \rightarrow \infty} \inf_{u \in \mathbb{H}_a} \mathbb{E} \left[-f(W_T + I_T(u)) + \Phi_T(W, u) + \lambda \int (I_T(u))^4 + \frac{1}{2} \|l^T(u)\|_{\mathcal{H}}^2 \right] \\ &= \inf_{u \in \mathbb{H}_a^{-1/2-\varepsilon}} \mathbb{E} \left[-f(W_{\infty} + I_{\infty}(u)) + \Phi_{\infty}(W, u) + \lambda \int (I_{\infty}(u))^4 + \frac{1}{2} \|l^{\infty}(u)\|_{\mathcal{H}}^2 \right] \end{aligned}$$

$$\int e^f d\Phi_{\frac{4}{3}} = e^{\mathcal{W}_{\infty}(0) - \mathcal{W}_{\infty}(f)} = \lim_{T \rightarrow \infty} \int e^f d\Phi_{\frac{4}{3}, T}.$$

▷ First “explicit” description of $\Phi_{\frac{4}{3}}$

Thanks.

