

...

Pathwise regularisation by noise in PDEs

...

Massimiliano Gubinelli – Université Paris Dauphine.

Workshop “Averaging and homogenisation in deterministic and stochastic systems” CIRM, Mai 2015.

- ▷ We will discuss some examples of the "good" effects of "bad", "irregular" functions.
- ▷ In particular we will look at non-linear differential (partial or ordinary) equations perturbed by some kind of (deterministic) noise.
- ▷ By defining a suitable notion of "irregular" noise we are able to show, in a quantitative way, that the more the noise is irregular the more the properties of the equation are better.
- ▷ Some examples includes: ODE perturbed by additive noise, linear stochastic transport equations and non-linear modulated dispersive PDEs.
- ▷ It is possible to show that the sample paths of Brownian motion or fractional Brownian motion and related processes have almost surely this kind of irregularity.

Joint work with Remi Catellier and Khalil Chouk.

Addition of noise has positive effects on the theory of the equation (in a pathwise sense)

→ ODEs:

$$X_t = x + \int_0^t b(X_s) ds + W_t$$

where (W_t) is a BM in \mathbb{R}^d and b a less-than-Lipshitz vectorfield. Many results: Veretenikov, Davie, Krylov-Röckner, Flandoli, Attanasio, Fedrizzi, Proske, ... Essentially: bounded b : (in L^∞ or with some particular integrability: LPS condition).

→ Transport equation:

$$d_t u(t, x) + b(x) \cdot \nabla u(t, x) dt = \nabla u(t, x) \cdot dW_t$$

good theory for L^∞ solutions and preservation of regularity. Flandoli-G.-Priola, Flandoli-Attanasio, Flandoli-Maurelli, Flandoli-Beck-G.-Maurelli

→ Some other PDE: Vlasov-Poisson, point vortices in 2d.

→ Modulated non-linear Schrödinger equation in $d=1$. De Bouard-Debussche, Debussche-Tsutsumi.

Goal: provide a deterministic framework to discuss regularization by “perturbations/modulation” for the following model PDEs:

- **Transport equation:** $x \in \mathbb{R}^d$, $t \geq 0$, $w: \mathbb{R} \rightarrow \mathbb{R}^d$, $b: \mathbb{R}^d \rightarrow \mathbb{R}^d$

$$\partial_t u(t, x) + \dot{w}_t \cdot \nabla u(t, x) + b(x) \cdot \nabla u(t, x) = 0, \quad u(0, \cdot) = u_0.$$

- **Non-linear Schrödinger equation:** $x \in \mathbb{T}, \mathbb{R}$, $t \geq 0$, $w: \mathbb{R} \rightarrow \mathbb{R}$

$$\partial_t \varphi(t, x) = i \Delta \varphi(t, x) \dot{w}_t + i |\varphi(t, x)|^{p-2} \varphi(t, x).$$

- **Korteweg–de Vries equation:** $x \in \mathbb{T}, \mathbb{R}$, $t \geq 0$, $w: \mathbb{R} \rightarrow \mathbb{R}$

$$\partial_t u(t, x) = \partial_x^3 u(t, x) \dot{w}_t + \partial_x (u(t, x))^2.$$

Consider the linear transport PDE

$$\partial_t u(t, x) + \dot{w}_t \cdot \nabla u(t, x) = f(x), \quad u(0, \cdot) = 0.$$

Solutions are given explicitly by

$$u(t, x) = \int_0^t f(x + w_s - w_t) ds = T_t^w f(x - w_t)$$

where given a function $w: [0, 1] \rightarrow \mathbb{R}^d$ we define the **averaging operator**

$$T_t^w f(x) = \int_0^t f(x + w_s) ds, \quad T_{t,s}^w f = T_t^w f - T_s^w f$$

acting on functions (or distributions) $f: \mathbb{R}^d \rightarrow \mathbb{R}$.

Question: What is the relation between w , the regularity of f and that of $u(t, \cdot)$?

If w is smooth we do not expect anything special to happen and u to have the same regularity of f .

▷ $d=1$, $w_t=t$. Then if $F'(x) = f(x)$ we have $T_t^w f(x) = \int_0^t F'(x+s)ds = F(x+t) - F(x)$ and $T^w: L^\infty \rightarrow \text{Lip}$:

$$|T_t^w f(x) - T_t^w f(y)| \leq \|f\|_\infty |x - y|, \quad |T_{t,s}^w f(x)| \leq \|f\|_\infty |t - s|$$

▷ Tao–Wright: if w “wiggles enough” then T_t^w maps L^q into $L^{q'}$ with $q' > q$.

▷ Davie: if w is a sample of BM then a.s. (the exceptional set depends on f)

$$|T_{t,s}^w f(x) - T_{t,s}^w f(y)| \leq C_w \|f\|_\infty |x - y|^{1-} |t - s|^{1/2-}$$

Problem: study the mapping properties of T^w for w the sample path of a stochastic process.

Consider

$$Y_t^w(\xi) = \int_0^t e^{i\langle \xi, w_s \rangle} ds$$

then $T_t^w f = \mathcal{F}^{-1}(Y_t^w \mathcal{F}(f))$. Mapping properties of T^w in $(H^s)_{s \in \mathbb{R}}$ spaces can be discussed in terms of Y^w :

$$\|T_{t,s}^w f\|_{H^s} = \|(1 + \xi^2)^{s/2} Y_{t,s}^w(\xi) \mathcal{F}f(\xi)\|_{H_\xi^s}.$$

In our setting more convenient to look at the scale $(\mathcal{FL}^\alpha)_\alpha$:

$$\|f\|_{\mathcal{FL}^\alpha} = \int |f(\xi)| (1 + \xi^2)^{\alpha/2} d\xi$$

since $C^\alpha \subseteq \mathcal{FL}^\alpha$.

Definition 1 (Catellier–G.) *We say that w is (ρ, γ) -irregular if there exists a constant K such that for all $\xi \in \mathbb{R}^d$ and $0 \leq s \leq t \leq 1$:*

$$|Y_{t,s}^w(\xi)| \leq K(1 + |\xi|)^{-\rho} |t - s|^\gamma.$$

Theorem 2 *The fBM of Hurst index H is ρ -irregular for any $\rho < 1/2H$.*

\Rightarrow there exists functions of arbitrarily high irregularity and arbitrarily L^∞ -near any given continuous function.

Lemma 3 *An irregular function cannot be too regular.*

Proof. If $w \in C^\theta$ with $\alpha\theta + \gamma > 1$ and $\alpha \in [0, 1]$, using the Young integral, we find

$$|t - s| = |e^{ia}(t - s)| = \left| \int_s^t \underbrace{e^{ia - iaw_r}}_{C^{\alpha\theta}} d_r \underbrace{Y_r^w(a)}_{C^\gamma} \right|$$

$$\leq C K_w (|t - s|^\gamma + |t - s|^{\alpha\theta + \gamma} |a|^\alpha) \|w\|_\theta (1 + |a|)^{-\rho} \rightarrow 0$$

if $t > s$ and $\alpha < \rho$. This implies that is not possible that $\theta > (1 - \gamma) / \rho$.

- ▷ Not easy to say if a function is irregular.
- ▷ In $d = 1$ smooth functions are (ρ, γ) irregular for $\rho + \gamma = 1$. In particular if we insist on $\gamma > 1/2$ we have $\rho < 1/2$.
- ▷ For $d > 1$ smooth functions are not irregular: if $|t - s| \ll 1$

$$\int_s^t e^{i\langle a, w_r \rangle} dr \simeq \int_s^t e^{i\langle a, w'_s \rangle (t-s)} dr \simeq (1 + |\langle a, w'_s \rangle|)^{-1} \not\sim (1 + |a|)^{-\rho}.$$

- ▷ If w is ρ -irregular and φ is a C^1 perturbation then $w + \varphi$ is at least $\rho - (1 - \gamma)$ irregular since:

$$Y_{t,s}^{w+\varphi}(\xi) = \int_s^t e^{i\langle \xi, w_r + \varphi_r \rangle} dr = \int_s^t e^{i\langle \xi, \varphi_r \rangle} dr Y_{s,r}^w(\xi)$$

and we can use Young integral estimates.

- ▷ If W is a fBM and Φ an adapted smooth perturbation then $W + \Phi$ is as irregular as W (via Girsanov theorem).
- ▷ Other results (see Catellier thesis): relation with intersection local times, irregularity for α -stable Levy processes, relation with local non-determinism.

Theorem 4 *If w is ρ -irregular then*

$$T^w: H^s \rightarrow H^{s+\rho}$$

and

$$T^w: \mathcal{FL}^\alpha \rightarrow \mathcal{FL}^{\alpha+\rho}.$$

Proof. Indeed

$$\begin{aligned} \|T_{t,s}^w f\|_{\mathcal{FL}^{\alpha+\rho}} &= \int d\xi (1 + |\xi|)^{\alpha+\rho} |Y_{t,s}^w(\xi)(\mathcal{F}f)(\xi)| \\ &\leq K_w |t - s|^\gamma \int d\xi (1 + |\xi|)^\alpha |(\mathcal{F}f)(\xi)| = K_w |t - s|^\gamma \|f\|_{\mathcal{FL}^\alpha}. \end{aligned}$$

Remark 5 More difficult to understand the mapping properties in other spaces, for example Hölder spaces C^α . Only partial results available.

- ▷ Consider the transport equation with a perturbation:

$$\partial_t u(t, x) + \dot{w}_t \cdot \nabla u(t, x) + b(x) \cdot \nabla u(t, x) = 0, \quad u(0, \cdot) = u_0.$$

- ▷ In the Lipschitz case there is only one solution u given by the method of characteristics:

$$u(t, x) = u_0(\phi_t^{-1}(x))$$

where $\phi_t(x) = x_t$ is the flow of the ODE

$$\begin{cases} \dot{x}_t = b(x_t) + \dot{w}_t \\ x_0 = x \end{cases}$$

- ▷ Uniqueness of solutions is related to the uniqueness (and smoothness) theory of the flow.

In order to exploit the averaging properties of w in the study of the ODE

$$x_t = x_0 + \int_0^t b(x_s) ds + w_t$$

we rewrite it in order to make the action of the averaging operator explicit: let $\theta_t = x_t - w_t$:

$$\theta_t = \theta_0 + \int_0^t b(w_s + \theta_s) ds = \theta_0 + \int_0^t (d_s G_s)(\theta_s)$$

where $G_s(x) = T_s^w b(x)$ so that $d_s G_s(x) = f(w_s + x)$.

If we assume that G is C^γ in time ($\gamma > 1/2$) with values in a space of regular enough functions we can study this equation as a Young type equation for $\theta \in C^\gamma$.

▷ **Non-linear Young integral:**

$$\int_0^t (d_s G_s)(\theta_s) = \lim_{\Pi} \sum_i G_{t_{i+1}, t_i}(\theta_{t_i})$$

This limit exists if $\theta \in C_t^\gamma$ and $G \in C_t^\gamma C_x^\nu$ with $\gamma(1 + \nu) > 1$. The integral is in C_t^γ .

Theorem 6 *The integral equation*

$$\theta_t = \theta_0 + \int_0^t (d_s G_s)(\theta_s)$$

is well defined for $\theta \in C^\gamma$ and $G \in C_t^\gamma C_{x,\text{loc}}^\nu$ with $(1 + \nu)\gamma > 1$.

- Existence of global solutions if G of linear growth.
- Uniqueness if $G \in C_t^\gamma C_{x,\text{loc}}^{\nu+1}$ and differentiable flow.
- Smooth flow if $G \in C_t^\gamma C_x^{\nu+k}$.

Theorem 7 *The equation*

$$x_t = x_0 + \int_0^t b(x_s) ds + w_t$$

has a unique solution for w ρ -irregular and $b \in \mathcal{FL}^\alpha$ for $\alpha > 1 - \rho$. In this case we can take $\theta \in C^1$ above and the condition for uniqueness (and Lipschitz flow) is $G \in C_t^\gamma C_x^{3/2}$.

▷ Say that x is controlled by w if $\theta = x - w \in C^\gamma$. In this case we have

$$I_x(b) = \int_0^t b(x_s) ds = \int_0^t (d_s T_s^w b)(\theta_s)$$

and the r.h.s. is well defined as soon as $T^w b \in C_t^\gamma C_x^\nu$.

▷ If w is ρ irregular and $b \in \mathcal{FL}^\alpha$ then $T^w b \in C_t^\gamma \mathcal{FL}_x^{\alpha+\rho}$ so if $\alpha + \rho \geq \nu$ we have $T^w b \in C_t^\gamma C_x^\nu$.

In this case $I_x(b)$ can be extended by continuity to all $b \in \mathcal{FL}^\alpha$ and in particular we have given a meaning to

$$\int_0^t b(x_s) ds$$


when b is a distribution *provided* x is controlled by a ρ -irregular path.

▷ For controlled paths the ODE

$$x_t = x_0 + \int_0^t b(x_s) ds + w_t$$

make sense even for certain distributions b as a Young equation for θ .

Regularization of ODEs at a glance



(work of R. Catellier)

We want to give a meaning and study the uniqueness issue for the transport equation

$$(\partial_t + b(x) \cdot \nabla + \dot{w}_t \cdot \nabla)u(t, x) = 0$$

for $u \in L^\infty$ and $w \in C^\sigma$ with $\sigma > 1/3$ such that (w, \mathbb{W}) is a geometric σ -Hölder rough path such that w is ρ -irregular. For the moment only in the case $\operatorname{div} b = 0$.

▷ **Weak formulation:** We consider u as a distribution: $u_t(\varphi) = \int dx \varphi(x) u(t, x)$ for all $\varphi \in L^1(\mathbb{R}^d)$. The integral formulation of the equation is

$$u_t(\varphi) - u_s(\varphi) = \int_s^t u_r(\nabla \cdot (b\varphi)) dr + \int_s^t u_r(\nabla \varphi) d_r w_r$$

for all $\varphi \in \mathcal{S}(\mathbb{R}^d)$ and $0 \leq s \leq t$.

We need to give a meaning to such an integral equation in order to discuss the regularization by noise phenomenon. (No way out!)

▷ It is possible via the theory of **controlled rough paths**.

Let (X, \mathbb{X}) be a σ -Hölder rough path with $\sigma > 1/3$:

$$\mathbb{X}_{t,s} = \mathbb{X}_{t,u} + \mathbb{X}_{u,s} + (X_t - X_u) \otimes (X_u - X_s), \quad |X_t - X_s| + |\mathbb{X}_{s,t}|^{1/2} = O(|t - s|^\sigma)$$

▷ We say that $y \in C_t^\sigma$ is **controlled by X** if there exists $y^X \in C_t^\sigma$ such that

$$y_t - y_s - y_s^X (X_t - X_s) =: y_{s,t}^\# = O(|t - s|^{2\sigma}).$$

▷ For a controlled path y we can define the integral against X by compensated Riemman sums:

$$I_t = \int_0^t y_s dX_s := \lim_{\Pi} \sum_i y_{t_i} (X_{t_{i+1}} - X_{t_i}) + y_{t_i}^X \mathbb{X}_{t_{i+1}, t_i}$$

▷ This integral is the only function (up to constants) which has the following property

$$I_t - I_s = y_s (X_t - X_s) + y_s^X \mathbb{X}_{t,s} + O(|t - s|^{3\sigma}).$$

In particular, the integral is itself controlled by X and $I^X = y$.

Definition 8 We say that u is a function controlled by w if for all $\varphi \in \mathcal{S}(\mathbb{R}^d)$ we have

$$u_t(\varphi) - u_s(\varphi) = u_s^w(\varphi)(w_t - w_s) + u_{t,s}^\sharp(\varphi)$$

where $u_s^w(\varphi) \in C^\sigma$ and $|u_{t,s}^\sharp(\varphi)| \lesssim |t - s|^{2\sigma}$.

Definition 9 If u is controlled we say that it is a L^∞ solution of the rough transport equation (RTE) if

$$u_t(\varphi) - u_s(\varphi) = \int_s^t u_r(\nabla \cdot (b\varphi))dr + \int_s^t u_r(\nabla \varphi)d_r w_r$$

holds for all $\varphi \in \mathcal{S}(\mathbb{R}^d)$, $0 \leq s \leq t$.

Remark: If $\sigma > 1/2$ we can just assume that $u_t(\nabla \varphi) \in C_t^\sigma$ so that the rough integral becomes a Young integral.

Equivalently, u is a solution to the RTE iff

$$u_t(\varphi) - u_s(\varphi) = \int_s^t u_r(\nabla \cdot (b\varphi))dr + u_s(\nabla \varphi)(w_t - w_s) + u_s(\nabla^2 \varphi)\mathbb{W}_{t,s} + O(|t - s|^{3\sigma})$$

Lemma 10 *If b is Lipschitz there exists a solution to the RTE given by $u(t, x) = u_0(\phi_t^{-1}(x))$.*

Proof. The proof proceed by approximation of (w, \mathbb{W}) by $(w^\varepsilon, \mathbb{W}^\varepsilon)$ and by stability of the flow. Let ϕ^ε be the approximate flow, then $u_t^\varepsilon(\varphi) = \int_{\mathbb{R}^d} u_0(\phi_t^{\varepsilon,-1}(x))\varphi(x)dx = \int_{\mathbb{R}^d} u_0(x)\varphi(\phi_t^\varepsilon(y))dy$. Taylor expansion gives

$$\varphi(\phi_t^\varepsilon(y)) = \varphi(\phi_s^\varepsilon(y)) + \int_s^t \nabla \varphi(\phi_r^\varepsilon(y))b(\phi_r^\varepsilon(y))dr + \nabla \varphi(\phi_s^\varepsilon(y))(w_t^\varepsilon - w_s^\varepsilon) + O_\varphi(|t - s|^{2\sigma})$$

That is $u_t^\varepsilon(\varphi) = u_s^\varepsilon(\varphi) + u_s^\varepsilon(\nabla \varphi)(w_t^\varepsilon - w_s^\varepsilon) + O_\varphi(|t - s|^{2\sigma})$. By weak compactness it is possible to pass to the limit (along a subsequence) in this equation and obtain a controlled path $u = \lim_{\varepsilon_k} u_{\varepsilon_k}$.

Uniqueness is proven by showing via a direct computation that

$$t \mapsto \int_{\mathbb{R}^d} u(t, \phi_t(x))\rho(x)dx = u_t(\rho \circ \phi_t^{-1})$$

is a constant function of t for all $\rho \in \mathcal{S}(\mathbb{R}^d)$. This implies that $u(t, \phi_t(x)) = u_0(x)$. Uniqueness depends only on the Lipschitz property of the flow.

Theorem 11 *Let $b \in \mathcal{FL}^\alpha$ for $\alpha > 0$ and $\alpha + \rho > 3/2$ and let w be ρ -irregular. Then there exists a unique solution to the RTE given by the method of characteristics.*

Proof. Approximate b by b_ε , then by the previous theorem there exists a unique solution u_ε to the RTE. Analysis of the approximate flow ϕ_ε shows that this solution converges to a controlled solution u of the RTE with vectorfield b . Since ϕ is Lipschitz we can prove again uniqueness. \square

Remark 12 The above result is path-wise. In particular b can depend on w .

Remark 13 If $b \in C^\alpha$, b deterministic and w is a fBm of Hurst index H then the uniqueness holds almost surely when $\alpha > 1 - 1/(2H)$ and $\alpha > 0$. This recovers the results of Flandoli–Gubinelli–Priola for the Brownian case but extend them well beyond the Brownian context.

(joint work with K. Chouk)

Two simple dispersive models with ρ -irregular modulation w :

- **Non-linear Schrödinger equation:** $x \in \mathbb{T}, \mathbb{R}, \mathbb{R}^2, t \geq 0$

$$\partial_t \varphi(t, x) = i\Delta \varphi(t, x) \partial_t w_t + i|\varphi(t, x)|^{p-2} \varphi(t, x).$$

- **Korteweg–de Vries equation:** $x \in \mathbb{T}, \mathbb{R}, t \geq 0$

$$\partial_t u(t, x) = \partial_x^3 u(t, x) \partial_t w_t + \partial_x (u(t, x))^2.$$

To be compared to the non-modulated setting where $\partial_t w_t = 1$ and studied in the scale of $(H^s)_s$ spaces.

The equations are understood in the mild formulation

$$u(t) = U_t^w u(0) + \int_0^t U_t^w (U_s^w)^{-1} \partial_x (u(s))^2 ds.$$

with $U_t^w = e^{i w_t \partial_x^3}$. (similarly for NLS). Here w can be an arbitrary continuous function.

Rewrite the mild formulation as $(U_t^w = e^{\partial_x^3 w t})$

$$v(t) = (U_t^w)^{-1}u(t) = u(0) + \int_0^t (U_s^w)^{-1} \partial_x (U_s^w v(s))^2 ds.$$

Theorem 14 *Let*

$$X_t(\varphi) = X_t(\varphi, \varphi) = \int_0^t (U_s^w)^{-1} \partial_x (U_s^w \varphi)^2 ds$$

If w is ρ irregular then $X \in C^\gamma \text{Lip}_{\text{loc}}(H^\alpha)$ for $\alpha > -\rho$ and $\rho > 3/4$.

For $v \in C^\gamma H^\alpha$ we can give a meaning to the non-linearity as a Young integral

$$\int_0^t (U_s^w)^{-1} \partial_x (U_s^w v(s))^2 ds := \int_0^t (d_s X_s)(v(s)) := \lim_{\Pi} \sum_i X_{t_{i+1}}(v(t_i)) - X_{t_i}(v(t_i))$$

The continuity of the Young integral implies that if $v_n \rightarrow v$ in $C^\gamma H^\alpha$ then

$$\int_0^t (U_s^w)^{-1} \partial_x (U_s^w v(s))^2 ds = \lim_n \int_0^t (U_s^w)^{-1} \partial_x (U_s^w v_n(s))^2 ds$$

Theorem 15 *The Young equation for $v \in C^\gamma H^\alpha$:*

$$v(t) = u(0) + \int_0^t (d_s X_s)(v(s))$$

has local solutions for initial conditions in H^α with locally Lipschitz flow. Uniqueness in $C^\gamma H^\alpha$.

▷ Equivalent “differential” formulation:

$$v(t) - v(s) = X_{t,s}(v(s)) + O(|t - s|^{2\gamma}), \quad v(0) = u_0$$

Regularization by modulation. In the non-modulated case it is known that there cannot be continuous flow for $\alpha \leq -1/2$ on \mathbb{T} and $\alpha \leq -3/4$ on \mathbb{R} .

- ▷ Global solutions thanks to the L^2 conservation and smoothing for $\alpha > 0$ or an adaptation of the I-method for $-3/2 \leq \alpha < 0$ and $\alpha > -\rho/(3 - 2\gamma)$.
- ▷ **NLS:** 1d, global solutions for $\alpha \geq 0$ and $\rho > 1/2$. 2d, local solutions for $\alpha \geq 1/2$.
- ▷ Global solutions for 1d NLS with $\alpha > 0$ come from a smoothing effect of the non-linearity which is due to the irregularity of the driving function.

A different line of attack to the modulated Schrödinger equation comes from the application of the following Strichartz type estimate which can be proved under the same ρ -irregularity assumption.

Theorem 16 *Let $T > 0$, $p \in (2, 5]$, $\rho > \min(\frac{3}{2} - \frac{2}{p}, 1)$ then there exists a finite constant $C_{w,T} > 0$ and $\gamma^*(p) > 0$ such that the following inequality holds:*

$$\left\| \int_0^\cdot U^\cdot(U_s^w)^{-1} \psi_s ds \right\|_{L^p([0,T], L^{2p}(\mathbb{R}))} \leq C_w T^{\gamma^*(p)} \|\psi\|_{L^1([0,T], L^2(\mathbb{R}))}$$

for all $\psi \in L^1([0, T], L^2(\mathbb{R}))$.

▷ In the deterministic case the Strichartz estimate does not have the factor of T in the critical case $p = 5$. This is a sign of a mild regularization effect of the noise.

As an application we obtain global well-posedness for the modulated NLS equation with generic power nonlinearity $i e: \mathcal{N}(\phi) = |\phi|^\mu \phi$: (Debussche–de Bouard, Debussche–Tsutsumi)

Theorem 17 *Let $\mu \in (1, 4]$, $p = \mu + 1$, $\rho > \min(1, 3/2 - \frac{2}{p})$ and $u^0 \in L^2(\mathbb{R})$ then there exists $T^* > 0$ and a unique $u \in L^p([0, T], L^{2p}(\mathbb{R}))$ such that the following equality holds:*

$$u_t = U_t^w u^0 + i \int_0^t U_t^w (U_s^w)^{-1} (|u_s|^\mu u_s) ds$$

for all $t \in [0, T^]$. Moreover we have that $\|u_t\|_{L^2(\mathbb{R})} = \|u_0\|_{L^2(\mathbb{R})}$ and then we have a global unique solution $u \in L_{loc}^p([0, +\infty), L^{2p}(\mathbb{R}))$ and $u \in C([0, +\infty), L^2(\mathbb{R}))$. If $u^0 \in H^1(\mathbb{R})$ then $u \in C([0, \infty), H^1(\mathbb{R}))$.*

Thanks.