

KPZ equation via paracontrolled distributions




Massimiliano Gubinelli

CEREMADE
Université Paris Dauphine

Kyoto University – October 24th, 2014



Fluctuations of a growing interface



A model for random interface growth (think e.g. expansion of colony of bacteria): $h: \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$,

$$\partial_t h(t, x) = \underbrace{\kappa \Delta h(t, x)}_{\text{relaxation}} + \underbrace{F(\partial_x h(t, x))}_{\text{slope-dependent growth}} + \underbrace{\eta(t, x)}_{\text{noise with microscopic correlations}}$$

Fluctuations of a growing interface



The Kardar–Parisi–Zhang equation

- ▶ Kardar–Parisi–Zhang '84: slope-dependent growth given by $F(\partial_x h)$, in a certain scaling regime of small gradients:

$$F(\partial_x h) = F(0) + F'(0)\partial_x h + F''(0)(\partial_x h)^2 + \dots$$

- ▶ KPZ equation is the **universal model** for random interface growth

$$\partial_t h(t, x) = \underbrace{\kappa \Delta h(t, x)}_{\text{relaxation}} + \underbrace{\lambda [(\partial_x h(t, x))^2 - \infty]}_{\text{renormalized growth}} + \underbrace{\xi(t, x)}_{\text{space-time white noise}}$$

- ▶ This derivation is **highly problematic** since $\partial_x h$ is a distribution. But: [Hairer, Quastel \(2014, unpublished\)](#) justify it rigorously via scaling of smooth models and small gradients.
- ▶ KPZ equation is suspected to be universal scaling limit for random interface growth models, random polymers, and many particle systems;
- ▶ contrary to Brownian setting: KPZ has **fluctuations of order $t^{1/3}$** ; large time limit distribution of $t^{-1/3}h(t, t^{2/3}x)$ is expected to be universal in a sense comparable only to the Gaussian distribution.

KPZ and its siblings:

- ▶ KPZ equation:

$$\mathcal{L}h(t, x) = "(\partial_x h(t, x))^2 - \infty" + \xi(t, x);$$

$h: \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$, $\mathcal{L} = \partial_t - \Delta$ heat operator, ξ , space-time white noise;

- ▶ Burgers equation:

$$\mathcal{L}u(t, x) = "\partial_x(u(t, x)^2)" + \partial_x \xi(t, x);$$

solution is (formally) given by derivative of the KPZ equation: $u = \partial_x h$;

- ▶ solution to KPZ (formally) given by Cole-Hopf transform of the **stochastic heat equation**: $h = \log w$, where w solves

$$\mathcal{L}w(t, x) = "w(t, x) \diamond \xi(t, x)".$$

- ▶ All three are **universal objects**, that are expected to be scaling limits of a wide range of particle systems.

Stochastic Burgers equation

Take $u = Dh$

$$\mathcal{L}u = D\xi + Du^2$$

to obtain the stochastic Burgers equation (SBE) with additive noise.

▷ **Invariant measure:** Formally the SBE leaves invariant the space white noise: if u_0 has a Gaussian distribution with covariance $\mathbb{E}[u_0(x)u_0(y)] = \delta(x - y)$ then for all $t \geq 0$ the random function $u(t, \cdot)$ has a Gaussian law with the same covariance.

▷ **First order approximation:** Let $X(t, x)$ be the solution of the linear equation

$$\partial_t X(t, x) = \partial_x^2 X(t, x) + \partial_x \xi(t, x), \quad x \in \mathbb{T}, t \geq 0$$

X is a stationary Gaussian process with covariance

$$\mathbb{E}[X(t, x)X(s, y)] = p_{|t-s|}(x - y).$$

Almost surely $X(t, \cdot) \in \mathcal{C}^\gamma$ for any $\gamma < -1/2$ and any $t \in \mathbb{R}$. For any $t \in \mathbb{R}$ $X(t, \cdot)$ has the law of the white noise over \mathbb{T} .

Littlewood-Paley blocks and Hölder-Besov spaces

We will measure regularity in Hölder-Besov spaces $\mathcal{C}^\gamma = B_{\infty,\infty}^\gamma(\mathbb{T}^d)$.
If time is involved we abuse the notation by setting $\mathcal{C}^\gamma = C([0, T], B_{\infty,\infty}^\gamma(\mathbb{T}^d))$.

$f \in \mathcal{C}^\gamma, \gamma \in \mathbb{R}$ iff

$$\|\Delta_i f\|_{L^\infty} \leq \|f\|_\gamma 2^{-i\gamma}, \quad i \geq -1.$$

$$\mathcal{F}(\Delta_i f)(\xi) = \rho_i(\xi) \hat{f}(\xi)$$

where $\rho_i : \mathbb{R}^d \rightarrow \mathbb{R}_+$ are smooth functions with support $\simeq 2^i \mathcal{A}$ when $i \geq 0$
and form a partition of unity $\sum_{i \geq -1} \rho_i(\xi) = 1$ for all $\xi \neq 0$ so that

$$f = \sum_{i \geq -1} \Delta_i f$$

in \mathcal{S}' .

Paraproducts

Deconstruction of a product: $f \in \mathcal{C}^\rho, g \in \mathcal{C}^\gamma$

$$fg = \sum_{i,j \geq -1} \Delta_i f \Delta_j g = f \prec g + f \circ g + f \succ g$$

$$f \prec g = g \succ f = \sum_{i < j-1} \Delta_i f \Delta_j g \quad f \circ g = \sum_{|i-j| \leq 1} \Delta_i f \Delta_j g$$

Paraproduct (Bony, Meyer et al.)

$$f \prec g \in \mathcal{C}^{\min(\gamma+\rho, \gamma)}$$
$$f \circ g \in \mathcal{C}^{\gamma+\rho} \quad \text{only if } \gamma + \rho > 0$$

Proof. Recall $f \in \mathcal{C}^\rho, g \in \mathcal{C}^\gamma$.

$$i \ll j \Rightarrow \text{supp } \mathcal{F}(\Delta_i f \Delta_j g) \subseteq 2^i \mathcal{A} \quad i \sim j \Rightarrow \text{supp } \mathcal{F}(\Delta_i f \Delta_j g) \subseteq 2^j \mathcal{B}$$

So if $\rho > 0$

$$\Delta_q(f \prec g) = \sum_{j:j \sim q} \sum_{i:i < j-1} \underbrace{\Delta_q(\Delta_i f \Delta_j g)}_{O(2^{-i\rho-j\gamma})} = O(2^{-q\gamma}) \Rightarrow f \prec g \in \mathcal{C}^\gamma,$$

while if $\rho < 0$

$$\Delta_q(f \prec g) = \sum_{j:j \sim q} \sum_{i:i < j-1} \underbrace{\Delta_q(\Delta_i f \Delta_j g)}_{O(2^{-i\rho-j\gamma})} = O(2^{-q(\gamma+\rho)}) \Rightarrow f \prec g \in \mathcal{C}^{\gamma+\rho}.$$

Finally for the resonant term we have

$$\Delta_q(f \circ g) = \sum_{i \sim j \geq q} \Delta_q(\Delta_i f \Delta_j g) = \sum_{i \geq q} O(2^{-j(\rho+\gamma)}) \Rightarrow f \circ g \in \mathcal{C}^{\gamma+\rho}$$

but *only if* the sum converges.

Small detour : Young integral

Take $f \in \mathcal{C}^\rho, g \in \mathcal{C}^\gamma$ with $\gamma, \rho \in (0, 1)$

$$fDg = \underbrace{f \prec Dg}_{\mathcal{C}^{\gamma-1}} + \underbrace{f \circ Dg + f \succ Dg}_{\mathcal{C}^{\gamma+\rho-1}}$$

then


$$\begin{aligned} \int fDg &= \int \underbrace{f \prec Dg}_{\mathcal{C}^\gamma} + \int \underbrace{(f \circ Dg + f \succ Dg)}_{\mathcal{C}^{\gamma+\rho}} \\ &= f \prec g + \mathcal{C}^{\gamma+\rho}. \end{aligned}$$

Compare with standard estimate for the Young integral in Hölder spaces (valid when $\gamma + \rho > 1$):

$$\int_s^t f_u dg_u = f_s(g_t - g_s) + O(|t - s|^{\gamma+\rho}).$$

Expansion in smallness of increments vs. Expansion in regularity

Paraproduct as frequency modulation



Some other problems in singular SPDEs /I

Define and solve (locally) the following SPDEs:

- ▶ Stochastic differential equations (1+0): $u \in [0, T] \rightarrow \mathbb{R}^n$

$$\partial_t u(t) = \sum_i f_i(u(t)) \xi^i(t)$$

with $\xi : \mathbb{R} \rightarrow \mathbb{R}^m$ m -dimensional white noise in time.

- ▶ Burgers equations (1+1): $u \in [0, T] \times \mathbb{T} \rightarrow \mathbb{R}^n$

$$\partial_t u(t, x) = \Delta u(t, x) + f(u(t, x)) Du(t, x) + \xi(t, x)$$

with $\xi : \mathbb{R} \times \mathbb{T} \rightarrow \mathbb{R}^n$ space-time white noise.

Recall that

$$\xi \in \mathcal{C}^{-d/2-}$$

Some other problems in singular SPDEs /II

- ▶ Generalized Parabolic Anderson model (1+2): $u \in [0, T] \times \mathbb{T}^2 \rightarrow \mathbb{R}$

$$\partial_t u(t, x) = \Delta u(t, x) + f(u(t, x))\xi(x)$$

with $\xi : \mathbb{T}^2 \rightarrow \mathbb{R}$ space white noise.

- ▶ Stochastic quantization equation (1+3)

$$\partial_t u(t, x) = \Delta u(t, x) + "u(t, x)^3" + \xi(t, x)$$

with $\xi : \mathbb{R} \times \mathbb{T}^3 \rightarrow \mathbb{R}$ space-time white noise.

- ▶ But (currently) not: Multiplicative SPDEs (1+1)

$$\partial_t u(t, x) = \Delta u(t, x) + f(u(t, x))\xi(t, x)$$

with $\xi : \mathbb{R} \times \mathbb{T} \rightarrow \mathbb{R}$ space-time white noise.

Joint work with P. Imkeller and N. Perkowski.

(Also K. Chouk and R. Catellier for $(\Phi)_3^4$).

The prototype of a singular PDE

Consider the simple controlled PDE (η smooth, fixed initial condition)

$$\partial_t u(t, x) = \Delta u(t, x) + F(u(t, x))\eta(x)$$

$u : \mathbb{R}_+ \times \mathbb{T}^d \rightarrow \mathbb{R}$, $\eta : \mathbb{T}^d \rightarrow \mathbb{R}$ and smooth function $F : \mathbb{R} \rightarrow \mathbb{R}$.

Problem

The solution map

$$\eta \xrightarrow{\Psi} u$$

is generally **not** continuous for $\eta \in \mathcal{C}^{\gamma-2}$ with $\gamma < 1$.

Reason: $u \in \mathcal{C}^\gamma$ and $\eta \in \mathcal{C}^{\gamma-2}$ cannot be multiplied when $2\gamma - 2 \leq 0$. The r.h.s. of the equation is not well defined.


What can go wrong?

Consider the sequence of functions $x^n : \mathbb{R} \rightarrow \mathbb{R}^2$

$$x(t) = \frac{1}{n}(\cos(2\pi n^2 t), \sin(2\pi n^2 t))$$

then $x^n(\cdot) \rightarrow 0$ in $\mathcal{C}^\gamma([0, T]; \mathbb{R}^2)$ for any $\gamma < 1/2$. But

$$I(x^{n,1}, x^{n,2})(t) = \int_0^t x^{n,1}(s) \partial_t x^{n,2}(s) ds \rightarrow \frac{t}{2} \neq I(0,0)(t) = 0$$



The definite integral $I(\cdot, \cdot)(t)$ is **not** a continuous map $\mathcal{C}^\gamma \times \mathcal{C}^\gamma \rightarrow \mathbb{R}$ for $\gamma < 1/2$.

(Cyclic microscopic processes can produce macroscopic results. Resonances.)

Homogeneisation of a random potential

▷ Consider the linear heat equation with a small random time-independent (Gaussian) potential V

$$\partial_t U(t, x) = \Delta U(t, x) + \varepsilon^{2-\alpha} V(x) U(t, x)$$

on $(\mathbb{T}/\varepsilon)^d$ and where ε is a small parameter and $\alpha < 2$.

▷ Introduce macroscopic variables $u_\varepsilon(t, x) = U(t/\varepsilon^2, x/\varepsilon)$ with parabolic rescaling, then

$$\partial_t u_\varepsilon(t, x) = \Delta u_\varepsilon(t, x) + V_\varepsilon(x) u_\varepsilon(t, x)$$

on \mathbb{T} and where $V_\varepsilon(x) = \varepsilon^{-\alpha} V(x/\varepsilon)$.

Homogeneisation of a random potential (II)

The covariance of the macroscopic noise is

$$\mathbb{E}[V_\varepsilon(x)V_\varepsilon(y)] = \varepsilon^{-2\alpha}C((x-y)/\varepsilon)$$

Theorem

If $d > 2\alpha$ then $V_\varepsilon \rightarrow 0$ in $\mathcal{C}^{-\alpha-}$. While if $d = 2\alpha$ then V_ε converges to the space white noise on \mathbb{T} .

So we are let to the study of the stability properites of the equation

$$\mathcal{L}u = \eta u$$

with $\eta \in \mathcal{C}^{-\alpha}$. This stability is easy to establish when $2 - 2\alpha > 0$ by standard estimates in Besov spaces. We are concerned then with the case $\alpha = 1$.

Transformation of PAM

▷ In order to understand the difficulties, let us perform a change of variable by letting $u = e^X v$ with $\mathcal{L}X = \eta$. We get

$$\begin{aligned}\mathcal{L}u &= v\mathcal{L}e^X + e^X\mathcal{L}v - \partial_x e^X \partial_x v \\ &= ve^X\mathcal{L}X - ve^X(\partial_x X)^2 + e^X\mathcal{L}v - e^X\partial_x X\partial_x v\end{aligned}$$

so v solves

$$\mathcal{L}v = (\partial_x X)^2 v + \partial_x X \partial_x v.$$

Let $\gamma = 2 - \alpha$ – the regularity of X .

▷ If we *assume* that $(\partial_x X)^2 \in \mathcal{C}^{2\gamma-2}$ then we see that this equation can be solved for $v \in \mathcal{C}^{2\gamma}$ since in this case $\partial_x X \partial_x v \in \mathcal{C}^{\gamma-1}$ and we have a continuous map

$$(X, (\partial_x X)^2) \in \mathcal{C}^\gamma \times \mathcal{C}^{2\gamma-2} \mapsto v \in \mathcal{C}^\gamma$$

Homogeneisation

When $\eta = V_\varepsilon$:

Theorem

Assume $d > 2$ and $\alpha = 1$ and let $\mathcal{L}X_\varepsilon = V_\varepsilon$ (+ technical conditions on the covariance C), then $(\partial_x X_\varepsilon)^2 \rightarrow \sigma^2$ in \mathcal{C}^{0-} .

▷ If $d > 2$ writing $u_\varepsilon = e^{X_\varepsilon} v_\varepsilon$ we obtain that v_ε converges to the solution of the PDE

$$\mathcal{L}v = \sigma^2 v$$

and so does u since $X \rightarrow 0$ in \mathcal{C}^γ .

▷ Now

$$\mathcal{L}u_\varepsilon = V_\varepsilon u_\varepsilon \not\rightarrow \mathcal{L}u = 0 * u$$

but $\mathcal{L}u = \sigma^2 u$ with $\sigma^2 \neq 0$. Lack of continuity of the problem wrt the data V_ε in the $\mathcal{C}^{\gamma-2}$ topology if $\gamma - 2 < -1$.

Renormalization

When $d = 2$, $\alpha = 1$:

Theorem

Let $\gamma = 1-$, then $V_\varepsilon \rightarrow \xi$ (white noise on \mathbb{T}^2) in $\mathcal{C}^{\gamma-2}$ and $\mathcal{L}X_\varepsilon = V_\varepsilon$ (+ technical conditions on the covariance C), then there exists a sequence $c_\varepsilon \rightarrow +\infty$ such that $(\partial_x X_\varepsilon)^2 - c_\varepsilon \rightarrow (\partial_x X)^{\circ 2}$ in $\mathcal{C}^{2\gamma-2}$.

Here, formally, $\sigma^2 = +\infty$, so there is not a well defined limit for u_ε .

Consider $\tilde{u}_\varepsilon(t, x) = e^{-c_\varepsilon t} u(t, x)$ which solves

$$\mathcal{L}\tilde{u}_\varepsilon = V_\varepsilon \tilde{u}_\varepsilon - c_\varepsilon \tilde{u}_\varepsilon$$

then for $\tilde{v}_\varepsilon = e^{-X_\varepsilon} \tilde{u}_\varepsilon$ we have the equation

$$\mathcal{L}\tilde{v}_\varepsilon = [(\partial_x X_\varepsilon)^2 - c_\varepsilon] \tilde{v}_\varepsilon + \partial_x X_\varepsilon \partial_x \tilde{v}_\varepsilon$$

which behaves well in the limit $\varepsilon \rightarrow 0$.

Structure of the explicit solution

▷ **Question:** What is the equation satisfied by $\tilde{u} = \lim_{\epsilon \rightarrow 0} \tilde{u}_\epsilon$?

It should be something like $\mathcal{L}\tilde{u} = \tilde{u}\xi - \infty\tilde{u} = \tilde{u} \diamond \xi$ (in which sense?)

▷ Note that (by parilinearization)

$$\tilde{u} = e^X \tilde{v} = e^X \prec \tilde{v} + e^X \succeq \tilde{v} = (e^X \prec X) \prec \tilde{v} + \mathcal{O}^{2\gamma} = \tilde{u} \prec X + \mathcal{O}^{2\gamma}$$

An analogous relation holds between \tilde{u}_ϵ and X_ϵ . Then

$$\begin{aligned} \tilde{u}_\epsilon V_\epsilon - c_\epsilon \tilde{u}_\epsilon &= \tilde{u}_\epsilon \prec V_\epsilon + \tilde{u}_\epsilon \circ V_\epsilon + \tilde{u}_\epsilon \succ V_\epsilon - c_\epsilon \tilde{u}_\epsilon \\ &= \tilde{u}_\epsilon \prec V_\epsilon + (\tilde{u}_\epsilon \prec X_\epsilon) \circ V_\epsilon + \tilde{u}_\epsilon^\sharp \circ V_\epsilon + \tilde{u}_\epsilon \succ V_\epsilon - c_\epsilon \tilde{u}_\epsilon \\ &= \tilde{u}_\epsilon \prec V_\epsilon + \tilde{u}_\epsilon (X_\epsilon \circ V_\epsilon - c_\epsilon) + C(\tilde{u}_\epsilon, X_\epsilon, V_\epsilon) + \tilde{u}_\epsilon^\sharp \circ V_\epsilon + \tilde{u}_\epsilon \succ V_\epsilon \end{aligned}$$

where we have used a **commutator lemma** which states roughly that

$$(\tilde{u}_\epsilon \prec X_\epsilon) \circ V_\epsilon \simeq \tilde{u}_\epsilon (X_\epsilon \circ V_\epsilon)$$

The main commutator estimate

All the difficulty is concentrated in the resonating term

$$f \circ g = \sum_{|i-j| \leq 1} \Delta_i f \Delta_j g$$

which however "is" smoother than $f \prec g$ if f or g has positive regularity.

Paraproducts decouple the problem from the source of the problem.

Commutator lemma

The trilinear operator $C(f, g, h) = (f \prec g) \circ h - f(g \circ h)$ satisfies

$$\|C(f, g, h)\|_{\beta+\gamma} \lesssim \|f\|_{\alpha} \|g\|_{\beta} \|h\|_{\gamma}$$

when $\beta + \gamma < 0$ and $\alpha + \beta + \gamma > 0$, $\alpha < 1$.

Structure of solution and paracontrolled distributions

▷ So in the limit $\varepsilon \rightarrow 0$ we have

$$\begin{aligned}\tilde{u}_\varepsilon V_\varepsilon - c_\varepsilon \tilde{u}_\varepsilon &= \tilde{u}_\varepsilon \prec V_\varepsilon + \tilde{u}_\varepsilon (X_\varepsilon \circ V_\varepsilon - c_\varepsilon) + C(\tilde{u}_\varepsilon, X_\varepsilon, V_\varepsilon) + \tilde{u}_\varepsilon^\sharp \circ V_\varepsilon + \tilde{u}_\varepsilon \succ V_\varepsilon \\ &\rightarrow \tilde{u} \prec \xi + \tilde{u}(X \diamond \xi) + C(\tilde{u}, X, \xi) + \tilde{u}^\sharp \circ \xi + \tilde{u} \succ \xi \\ &=: \tilde{u} \diamond \xi = \Phi(\tilde{u}, \tilde{u}^\sharp, X, X \diamond \xi)\end{aligned}$$

where $X \diamond \xi := \lim_{\varepsilon \rightarrow 0} (X_\varepsilon \circ V_\varepsilon - c_\varepsilon)$.

▷ **Question:** What is the equation satisfied by $\tilde{u} = \lim_{\varepsilon \rightarrow 0} \tilde{u}_\varepsilon$?

Indeed

$$\mathcal{L}\tilde{u} = \tilde{u} \diamond \xi = \Phi(\tilde{u}, \tilde{u}^\sharp, X, X \diamond \xi).$$

Where the r.h.s. is well defined since \tilde{u} is **paracontrolled** by X .

Paracontrolled distributions

Use the paraproduct to *define* a controlled structure. We say $y \in \mathcal{D}_x^\rho$ if $x \in \mathcal{C}^\gamma$

$$y = y^x \prec x + y^\sharp$$

with $y^x \in C^{\rho-\gamma}$ and $y^\sharp \in C^\rho$.

▷ **Paralinearization.** Let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be a sufficiently smooth function and $x \in \mathcal{C}^\gamma, \gamma > 0$. Then

$$\varphi(x) = \varphi'(x) \prec x + \mathcal{C}^{2\gamma}$$

▷ Another commutator: $f, g \in \mathcal{C}^{\rho-\gamma}, x \in \mathcal{C}^\gamma$

$$f \prec (g \prec h) = (fg) \prec h + \mathcal{C}^\rho$$

▷ **Stability.** ($\rho \leq 2\gamma$)

$$\varphi(y) = (\varphi'(y)y^x) \prec x + \mathcal{C}^\rho$$

so we can take $\varphi(y)^x = \varphi'(y)y^x$.

Solution theory for general signals

Goal: Show that $\Psi : \eta \mapsto u$ factorizes as

$$\eta \xrightarrow{J} J(\eta) \xrightarrow{\Phi} u$$

▷ *Analytic step:* show that when $\gamma \in (2/3, 1)$:

$$\Phi : \mathcal{X} \rightarrow \mathcal{C}^\gamma$$

is continuous. $\mathcal{X} = \overline{\text{Im}J} \subseteq \mathcal{C}^{\gamma-2} \times \mathcal{C}^{2\gamma-2}$ is the space of *enhanced signals* (or rough paths, or models).

But in general J is **not** a continuous map $\mathcal{C}^{\gamma-1} \rightarrow \mathcal{C}^{\gamma-2} \times \mathcal{C}^{2\gamma-2}$.

▷ *Probabilistic step:* prove that there exists a "reasonable definition" of $J(\xi)$ when ξ is a white noise. $J(\xi)$ is an explicit polynomial in ξ , so direct computations are possible.

Tools: Besov embeddings $L^p(\Omega; \mathcal{C}^\theta) \rightarrow L^p(\Omega; B_{p,p}^{\theta'}) \simeq B_{p,p}^{\theta'}(L^p(\Omega))$, Gaussian hypercontractivity $L^p(\Omega) \rightarrow L^2(\Omega)$, explicit L^2 computations.

Paracontrolled gPAM (I) - the r.h.s.

$u : \mathbb{R}_+ \hat{\mathbb{E}} \times \mathbb{T}^2 \rightarrow \mathbb{R}$, $\xi \in \mathcal{C}^{\gamma-2}$, $\gamma = 1-$. We want to solve (have uniform bounds for)

$$\mathcal{L}u = F(u)\xi = F(u) \prec \xi + F(u) \circ \xi + F(u) \succ \xi.$$

▷ Paracontrolled ansatz. Take $\mathcal{L}X = \xi$, $X \in \mathcal{C}^\gamma$ and assume that $u \in \mathcal{D}_X^{2\gamma}$:

$$u = u^X \prec X + u^\sharp$$

with $u^\sharp \in \mathcal{C}^{2\gamma}$ and $u^X \in \mathcal{C}^\gamma$.

▷ Paralinearization:

$$F(u) = F'(u) \prec u + \mathcal{C}^{2\gamma} = (F'(u)u^X) \prec X + \mathcal{C}^{2\gamma}$$

▷ Commutator lemma:

$$\begin{aligned} F(u) \circ \xi &= ((F'(u)u^X) \prec X) \circ \xi + \mathcal{C}^{2\gamma} \circ \xi \\ &= \underbrace{(F'(u)u^X)(X \circ \xi)}_{\in \mathcal{C}^{2\gamma-2}} + \underbrace{C(F'(u)u^X, X, \xi) + \mathcal{C}^{2\gamma} \circ \xi}_{\in \mathcal{C}^{3\gamma-2}} \end{aligned}$$

if we assume that $(X \circ \xi) \in \mathcal{C}^{2\gamma-2}$.

Paracontrolled gPAM (II) - the l.h.s.

So if u is paracontrolled by X :

$$u = u^X \prec X + u^\sharp$$

and if $X \circ \xi \in \mathcal{C}^{2\gamma-2}$ we have a control on the r.h.s. of the equation:

$$F(u)\xi = \underline{F(u)} \prec \xi + F'(u)u^X(X \circ \xi) + \mathcal{C}^{3\gamma-2}$$

What about the l.h.s.?

$$\mathcal{L}u = \mathcal{L}u^X \prec X + \underline{u^X} \prec \xi + \mathcal{L}u^\sharp - \partial_x u^X \prec \partial_x X$$

so letting $u^X = F(u)$ we have

$$\mathcal{L}u^\sharp = -\mathcal{L}F(u) \prec X + F'(u)F(u)(X \circ \xi) + \mathcal{C}^{2\gamma-2}$$

Paracontrolled gPAM (III) - the paracontrolled fixed point.

The PDE

$$\mathcal{L}u = F(u)\xi,$$

is equivalent to the system

$$\begin{aligned}\partial_t X &= \xi, \\ \partial_t u^\sharp &= (F'(u)F(u))(X \circ \xi) - \underbrace{\mathcal{L}f(u) \prec X}_{\in \mathcal{C}^{2\gamma-2}} + \underbrace{R(f, u, X, \xi)}_{\in \mathcal{C}^{3\gamma-2}} \circ \xi, \\ u &= F(u) \prec X + u^\sharp\end{aligned}$$

▷ The system can be solved by fixed point (for small time) in the space $\mathcal{D}_X^{2\gamma}$ if we assume that

$$X \in \mathcal{C}^\gamma, \quad (X \circ \xi) \in \mathcal{C}^{2\gamma-2}.$$

Structure of the paracontrolled solution

▷ When ξ smooth, the solution to

$$\partial_t u = F(u)\xi, \quad u(0) = u_0$$

is given by $u = \Phi(u_0, \xi, X \circ \xi)$ where

$$\Phi : \mathbb{R}^d \times \mathcal{C}^{\gamma-2} \times \mathcal{C}^{2\gamma-2} \rightarrow \mathcal{C}^\gamma$$

is continuous for any $\gamma > 2/3$ and $z = \Phi(u_0, \xi, \varphi)$ is given by

$$\begin{cases} z = F(z) \prec X + z^\sharp \\ \partial_t z^\sharp = (F'(z)F(z))\varphi - \underbrace{\mathcal{L}F(z) \prec X}_{\in \mathcal{C}^{2\gamma-2}} + \underbrace{R(F, z, X, \xi) \circ \xi}_{\in \mathcal{C}^{3\gamma-2}} \end{cases}$$

▷ If $(\xi^n, X^n \circ \xi^n) \rightarrow (\xi, \eta)$ in $\mathcal{C}^{\gamma-2} \times \mathcal{C}^{2\gamma-2}$ and

$$\partial_t u^n = f(u^n)\xi^n, \quad u(0) = u_0$$

then $u^n \rightarrow u = \Phi(u_0, \xi, \eta)$.

Relaxed form of the PDE

▷ Note that in general we can have $\xi^{1,n} \rightarrow \xi$, $\xi^{2,n} \rightarrow \xi$ and

$$\lim_n X^{1,n} \circ \xi^{1,n} \neq \lim_n X^{2,n} \circ \xi^{2,n}$$

▷ Take ξ^n, ξ smooth but $\xi^n \rightarrow \xi$ in $\mathcal{C}^{\gamma-2}$. It can happen that

$$\lim_n X^n \circ \xi^n = X \circ \xi + \varphi \in \mathcal{C}^{2\gamma-1}$$

In this case $u^n \rightarrow u$ and $u = \Phi(\xi, X \circ \xi + \varphi)$ solves the equation

$$\mathcal{L}u = F(u)\xi + F'(u)F(u)\varphi.$$

The limit procedure generates correction terms to the equation.

The original equation **relaxes** to another form in which additional terms are generated.

"Ito" form of the PDE

In the smooth setting $u = \Phi(\xi, X \circ \xi + \varphi)$ solves

$$\mathcal{L}u = F(u)\xi + F'(u)F(u)\varphi.$$

If we choose $\varphi = -X \circ \xi$, then

$$v = \Phi(\xi, X \circ \xi + \varphi) = \Phi(\xi, 0)$$

solves

$$\mathcal{L}v = F(v)\xi - F'(v)F(v)X \circ \xi$$

and has the particular property of being a continuous map of $\xi \in \mathcal{C}^{\gamma-2}$ alone.

The renormalization problem

If ξ is the space white noise we have

$$\xi \in \mathcal{C}^{-1-}, \quad X \in C([0, T]; \mathcal{C}^{1-})$$

and

$$\begin{aligned} X \circ \xi &= X \circ \mathcal{L}X = \frac{1}{2} \mathcal{L}(X \circ X) + \frac{1}{2} (DX \circ DX) \\ &= \frac{1}{2} \mathcal{L}(X \circ X) - (DX \prec DX) + \frac{1}{2} (DX)^2 \end{aligned}$$

But now

$$\frac{1}{2} (DX)^2 = c + C \mathcal{C}^{0-}$$

with $c = +\infty!$.

No obvious definition of $X \circ \xi$ can be given. But there exists c_ε such that

$$X_\varepsilon \circ \xi_\varepsilon - c_\varepsilon \rightarrow "X \diamond \xi" \quad \text{in } \mathcal{C}^{0-}.$$

The renormalized gPAM

To cure the problem we add a suitable counterterm to the equation

$$\mathcal{L}u = f(u) \diamond \xi = f(u)\xi - c(f'(u)f(u))$$

this defines a new product, denoted by \diamond . Now

$$f(u) \circ \xi - c(f'(u)f(u)) = (f'(u)f(u))(X \circ \xi - c) + C(f'(u)f(u), X, \xi) + R(f, u, X) \circ \xi$$

▷ The renormalized gPAM is equivalent to the equation

$$\begin{aligned} \mathcal{L}u^\sharp &= -\mathcal{L}f(u) \prec X + Df(u) \prec DX + (f'(u)f(u))(X \circ \xi - c) \\ &\quad + C(f'(u)f(u), X, \xi) + R(f, u, X) \circ \xi \end{aligned}$$

together with $u = f(u) \prec X + u^\sharp$ and where

$$X \in \mathcal{C}^{1-}, \quad X \diamond \xi = (X \circ \xi - c) \in \mathcal{C}^{0-}, \quad u^\sharp \in \mathcal{C}^{2-}.$$

Finally a theorem

Theorem

Let $d = 2$, $\alpha = 1$, $\gamma = 1$ – and small $T > 0$. There exist constants c_ε such that letting u_ε the solution to

$$\mathcal{L}u_\varepsilon = V_\varepsilon F(u_\varepsilon) - c_\varepsilon F'(u_\varepsilon)$$

then $u_\varepsilon \rightarrow u$ in \mathbb{C}^γ as $\varepsilon \rightarrow 0$ and $u \in \mathcal{D}_X^{2\gamma}$ is the unique weak solution in $\mathcal{D}_X^{2\gamma}$ to the equation

$$\mathcal{L}u = \xi \diamond F(u) = F(u) \prec \xi + F'(u)(X \diamond \xi) + G(u^X, u^\#, X)$$

where

$$\xi = \lim_{\varepsilon \rightarrow 0} V_\varepsilon, \quad X \diamond \xi = \lim_{\varepsilon \rightarrow 0} X_\varepsilon \circ V_\varepsilon - c_\varepsilon$$

in $\mathbb{C}^{\gamma-2}$ and $\mathbb{C}^{2\gamma-2}$ resp. and ξ has the law of the white noise on \mathbb{T}^2 .

Other applications

- ▶ [Gubinelli, Imkeller, P. \(2012\)](#): Multidimensional extension of [Hairer's \(2011\)](#) generalized Burgers equation ($\sigma - d/2 > 1/3$):

$$\partial_t u(t, x) = -(-\Delta)^\sigma u(t, x) + G(u(t, x))D_x u(t, x) + \xi(t, x);$$

- ▶ [Catellier, Chouk \(2013\)](#): Stochastic quantization equation ϕ_3^4 ($d = 3$):

$$\mathcal{L}u(t, x) = -u(t, x)^{\circ 3} + \xi(t, x);$$

- ▶ [Furlan \(2014\)](#): Stochastic Navier Stokes equation ($d = 3$):

$$\mathcal{L}u(t, x) = -P((u(t, x) \cdot \nabla)u(t, x)) + \xi(t, x).$$

Expansion for the SBE

Recall the SBE:

$$\mathcal{L}u = Du^2 + \xi$$

▷ Let $u = X + u_1$ then

$$\mathcal{L}u_1 = \partial_x(u_1 + X)^2 = \underbrace{\partial_x X^2}_{-2-} + 2\partial_x(u_1 X) + \partial_x u_1^2$$

▷ Let X^\vee be the solution to

$$\mathcal{L}X^\vee = \partial_x X^2 \quad \Rightarrow \quad X^\vee \in \mathcal{C}^{0-}$$

and decompose further $u_1 = X^\vee + u_2$. Then

$$\mathcal{L}u_2 = \underbrace{2\partial_x(X^\vee X)}_{-3/2-} + 2\partial_x(u_2 X) + \underbrace{\partial_x(X^\vee X^\vee)}_{-1-} + 2\partial_x(u_2 X^\vee) + \partial_x(u_2)^2$$

▷ Define $\mathcal{L}X^\vee = 2\partial_x(X^\vee X)$ and $u_2 = X^\vee + u_3$ then $X^\vee \in \mathcal{C}^{1/2-}$

$$\mathcal{L}u_3 = \underbrace{2\partial_x(u_3 X)}_{-3/2-} + \underbrace{2\partial_x(X^\vee X)}_{-3/2-} + \underbrace{\partial_x(X^\vee X^\vee)}_{-1-} + 2\partial_x(u_2 X^\vee) + \partial_x(u_2)^2$$

Expansion /II

▷ The partial expansion for the solution reads

$$u = X + X^{\vee} + 2X^{\forall} + U$$

$$\begin{aligned}\mathcal{L}U &= 2\partial_x(UX) + 2\partial_x(X^{\forall}X) + \partial_x(X^{\vee}X^{\vee}) + 2\partial_x((2X^{\forall} + U)X^{\vee}) + \partial_x(2X^{\forall} + U)^2 \\ &= 2\partial_x(UX) + \mathcal{L}(2X^{\forall} + X^{\forall}) + 2\partial_x((2X^{\forall} + U)X^{\vee}) + \partial_x(2X^{\forall} + U)^2\end{aligned}$$

and the regularities for the driving terms

X	X^{\vee}	X^{\forall}	$X^{\forall\forall}$	$X^{\forall\forall\forall}$
$-1/2-$	$0-$	$1/2-$	$1/2-$	$1-$

We can assume $U \in \mathcal{C}^{1/2-}$ so that the terms

$$2\partial_x((2X^{\forall} + U)X^{\vee}) + \partial_x(2X^{\forall} + U)^2$$

are well defined.

The remaining problem is to deal with $2\partial_x(UX)$.

Paracontrolled ansatz for SBE

▷ Make the following ansatz $U = U' \prec Q + U^\sharp$. Then

$$\mathcal{L}U = \mathcal{L}U' \prec Q + U' \prec \mathcal{L}Q - \partial_x U' \prec \partial_x Q + \mathcal{L}U^\sharp$$

while

$$\begin{aligned}\mathcal{L}U &= 2\partial_x(UX) + \underbrace{\mathcal{L}(2X^{\vee\vee} + X^{\vee\vee}) + 2\partial_x((2X^{\vee\vee} + U)X^{\vee\vee}) + \partial_x(2X^{\vee\vee} + U)^2}_{R(U)} \\ &= 2\partial_x(U \prec X) + 2\partial_x(U \circ X) + 2\partial_x(U \succ X) + R(U) \\ &= 2(U \prec \partial_x X) + 2(\partial_x U \prec X) + 2\partial_x(U \circ X) + 2\partial_x(U \succ X) + R(U)\end{aligned}$$

so we can set $U' = 2U$ and $\mathcal{L}Q = \partial_x X$ and get the equation

$$\mathcal{L}U^\sharp = -\mathcal{L}U' \prec Q + \partial_x U' \prec \partial_x Q + 2(\partial_x U \prec X) + 2\partial_x(U \circ X) + 2\partial_x(U \succ X) + R(U)$$

▷ Observe that $Q, U, U' \in \mathcal{C}^{1/2-}$ and we can assume that $U^\sharp \in \mathcal{C}^{1-}$.

Commutator

- ▷ The difficulty is now concentrated in the resonant term $U \circ X$ which is not well defined.
- ▷ The paracontrolled ansatz and the commutation lemma give

$$U \circ X = (2U \prec Q) \circ X + U^\sharp \circ X = 2U(Q \circ X) + \underbrace{C(2U, Q, X)}_{1/2-} + \underbrace{U^\sharp \circ X}_{1/2-}$$

- ▷ A stochastic estimate shows that $Q \circ X \in \mathcal{C}^{0-}$

Paracontrolled solution to SBE

▷ The final system reads

$$u = X + X^{\vee} + 2X^{\heartsuit} + U$$

$$U = U' \prec Q + U^{\sharp}, \quad U' = 2X^{\heartsuit} + 2U$$

$$\begin{aligned} \mathcal{L}U^{\sharp} = & 4\partial_x(U(Q \circ X)) + 4\partial_x C(U, Q, X) + 2\partial_x(U^{\sharp} \circ X) - 2\mathcal{L}U \prec Q \\ & + 2\partial_x U \prec \partial_x Q + 2(\partial_x U \prec X) + 2\partial_x(U \succ X) + R(U) \end{aligned}$$

▷ This equation has a (local in time) solution $U = \Phi(J(\xi))$ which is a continuous function of the data $J(\xi)$ given by a collection of multilinear functions of ξ :

$$J(\xi) = (X, X^{\vee}, X^{\heartsuit}, X^{\spadesuit}, X^{\clubsuit}, X \circ Q)$$

Burgers equation and paracontrolled distributions

$$\mathcal{L}u(t, x) = \partial_x u^2(t, x) + \partial_x \xi(t, x), \quad u(0) = u_0.$$

Paracontrolled Ansatz

$u \in \mathcal{P}_{\text{rbe}}$ if $u = X + X^\vee + 2X^\heartsuit + u^\mathcal{Q}$ with

$$u^\mathcal{Q} = u' \prec Q + u^\sharp.$$

- ▶ Paracontrolled structure: Can define u^2 continuously as long as $(Q \circ X) \in C([0, T], \mathcal{C}^{0-})$ is given (together with tree data $X, X^\vee, X^\heartsuit, X^\spadesuit, X^\clubsuit$).
- ▶ Obtain local existence and uniqueness of paracontrolled solutions. Solution depends pathwise continuously on extended data $J(\xi) = (\xi, X, X^\vee, X^\heartsuit, X^\spadesuit, X^\clubsuit, Q \circ X)$.

KPZ equation

KPZ equation:

$$\mathcal{L}h(t, x) = (\partial_x h(t, x))^2 + \xi(t, x), \quad h(0) = h_0.$$

Expect $h(t) \in \mathcal{C}^{1/2-}$, so $\partial_x h(t) \in \mathcal{C}^{-1/2-}$ and $(\partial_x h(t))^2$ not defined. But: expand

$$u = Y + Y^{\vee} + 2Y^{\vee\vee} + h^P,$$

where $\mathcal{L}Y = \xi$, $\mathcal{L}Y^{\vee} = \partial_x Y \partial_x Y, \dots$. In general: $\partial_x Y^{\tau} = X^{\tau}$. Make **paracontrolled ansatz** for h^P :

$$h^P = \pi_{<}(h', P) + h^{\sharp}$$

with $h' \in C([0, T], \mathcal{C}^{1/2-})$, $h^{\sharp} \in C([0, T], \mathcal{C}^{2-})$, $\mathcal{L}P = X$. Write $h \in \mathcal{P}_{\text{kpz}}$.

Can define $(\partial_x h(t))^2$ for $h \in \mathcal{P}_{\text{kpz}}$ and obtain local existence and uniqueness of solutions.

KPZ and Burgers equation

$h \in \mathcal{P}_{\text{kpz}}$ if

$$h = Y + Y^\vee + 2Y^\psi + h^P, \quad h^P = h' \prec P + h^\sharp.$$

$u \in \mathcal{P}_{\text{rbe}}$ if

$$u = X + X^\vee + 2X^\psi + u^Q, \quad u^Q = u' \prec Q + u^\sharp.$$

- ▶ If $h \in \mathcal{P}_{\text{kpz}}$, then $\partial_x h \in \mathcal{P}_{\text{rbe}}$.
- ▶ If h solves KPZ equation, then $u = \partial_x h$ solves Burgers equation with initial condition $u(0) = \partial_x h_0$.
- ▶ If $u \in \mathcal{P}_{\text{rbe}}$, then any solution h of $\mathcal{L}h = u^2 + \xi$ is in \mathcal{P}_{kpz} .
- ▶ If u solves Burgers equation with initial condition $u(0) = \partial_x h_0$, and h solves $\mathcal{L}h = u^2 + \xi$ with initial condition $h(0) = h_0$, then h solves KPZ equation.

KPZ and heat equation

Heat equation:

$$\mathcal{L}w(t, x) = w(t, x) \diamond \xi(t, x) = w(t, x)\xi(t, x) - w(t, x) \cdot \infty, \quad w(0) = w_0.$$

Paracontrolled ansatz: $w \in \mathcal{P}_{\text{rhe}}$ if

$$w = e^{Y+Y^\vee+2Y^\heartsuit} w^P, \quad w^P = \pi_{<}(w', P) + w^\sharp$$

(comes from Cole-Hopf transform).

- ▶ Slightly cheat to make sense of product $w \diamond \xi$ for $w \in \mathcal{P}_{\text{rhe}}$:

$$\begin{aligned} w \diamond \xi &= \mathcal{L}w - e^{Y+Y^\vee+2Y^\heartsuit} \left[\mathcal{L}w^P - [\mathcal{L}(Y^\vee + Y^\heartsuit) + (\partial_x(Y + Y^\vee + 2Y^\heartsuit))^2]w^P \right] \\ &\quad + 2e^{Y+Y^\vee+2Y^\heartsuit} \partial_x(Y + Y^\vee + 2Y^\heartsuit) \partial_x w^P; \end{aligned}$$

(agrees with renormalized pointwise product $w \diamond \xi$ in smooth case and with Itô integral in white noise case, continuous in extended data).

- ▶ Obtain global existence and uniqueness of solutions.
- ▶ One-to-one correspondence between \mathcal{P}_{kpz} and strictly positive elements of \mathcal{P}_{rhe} .
- ▶ Any solution of KPZ gives solution of heat equation. Any strictly positive solution of heat equation gives solution of KPZ equation.

Thanks

Para-modelled distributions

Let $\gamma > 0$ and (T, Π, Γ) regularity structure. Say f is **para-modelled**, $f \in \mathcal{P}^\gamma$, if there exists $f^\pi \in \mathcal{D}^\gamma$, with

$$f - \pi_{<}(f^\pi, \Pi) \in C^\gamma.$$

Example: $\mathcal{R}f^\pi \in \mathcal{P}^\gamma$.

Consider **rough path model**, say

$T = \text{span}(\Xi, \mathcal{I}(\Xi)\Xi, \mathcal{I}(\mathcal{I}(\Xi)\Xi)\Xi, \mathbf{1}, \mathcal{I}(\Xi), \mathcal{I}(\mathcal{I}(\Xi)\Xi))$. Try to solve

$\partial_t u = F(u)\xi$.

(Simplified) **para-modelled ansatz**: $u = \mathcal{R}u^\pi = \pi_{<}(u^\pi, \Pi) + u^\sharp$ with $u^\pi \in \mathcal{D}^{3\alpha}$.

Equation for u^\sharp :

$$\partial_t u^\sharp = -\partial_t \pi_{<}(u^\pi, \Pi) + F(u)\xi = \pi_{<}(u^\pi, D\Pi) - \pi_{<}(F(u^\pi) \star \xi^\pi, \Pi) + \text{smooth}.$$

To have $u^\sharp \in C^{3\alpha}$: choose expansion u^π so that all coefficients for terms of homogeneity $< 3\alpha - 1$ cancel. Obtain **a priori bounds** on $\|u^\sharp\|_{3\alpha}$ and then on $\|u^\pi\|_{\mathcal{D}^{3\alpha}}$. Thus at least **local existence** of solutions.

Stochastic Quantization

Stochastic quantization of $(\Phi^4)_3$: $\xi \in C^{-5/2-}$, $u \in C^{-1/2-}$, $u = u_1 + u_2 + u_{\geq 3}$.

$$\mathcal{L}u = \xi + \lambda(u^3 - 3c_1u - c_2u)$$

$$\mathcal{L}u_1 + \mathcal{L}u_{\geq 2} = \xi + \lambda(u_1^3 - 3c_1u_1) + 3\lambda(u_{\geq 2}(u_1^2 - c_1)) + 3\lambda(u_{\geq 2}^2u_1) + \lambda u_{\geq 2}^3 - \lambda c_2u$$

$$\triangleright \mathcal{L}u_1 = \xi \Rightarrow u_1 \in C^{-1/2-}, \mathcal{L}u_2 = \lambda(u_1^3 - 3c_1u_1) \Rightarrow u_2 \in C^{1/2-}$$

$$\mathcal{L}u_{\geq 3} = 3\lambda(u_{\geq 2}(u_1^2 - c_1)) + 3\lambda(u_2^2u_1) + 6\lambda(u_{\geq 3}u_2u_1) + 3\lambda(u_{\geq 3}^2u_1) + \lambda u_{\geq 3}^3 - \lambda c_2u$$

$$\triangleright \text{Ansatz: } u_{\geq 3} = 3\lambda u_{\geq 2} \prec X + u^\sharp, \text{ with } \mathcal{L}X = (u_1^2 - c_1)$$

$$\mathcal{L}u^\sharp = -3\lambda \mathcal{L}u_{\geq 2} \prec X + 3\lambda D u_{\geq 2} \prec DX + 3\lambda(u_{\geq 2} \circ (u_1^2 - c_1) - c_2u) + 3\lambda(u_{\geq 2} \succ (u_1^2 - c_1))$$

$$+ 3\lambda(u_2^2u_1) + 6\lambda(u_{\geq 3}(u_2u_1)) + 3\lambda(u_{\geq 3}^2u_1) + \lambda u_{\geq 3}^3$$

$$u_{\geq 2} \circ (u_1^2 - c_1) - c_2u = (u_2 \circ (u_1^2 - c_1) - c_2u_1) + (u_{\geq 3} \circ (u_1^2 - c_1) - c_2u_{\geq 2})$$

$$(u_{\geq 3} \circ (u_1^2 - c_1) - c_2u_{\geq 2}) = (3\lambda(u_{\geq 2} \prec X) \circ (u_1^2 - c_1) - c_2u_{\geq 2}) + u^\sharp \circ (u_1^2 - c_1)$$

$$= u_{\geq 2}(3\lambda(X \circ (u_1^2 - c_1)) - c_2) + 3\lambda \mathcal{C}(u_{\geq 2}, X, (u_1^2 - c_1)) + u^\sharp \circ (u_1^2 - c_1)$$

$$\triangleright \text{Basic objects: } (u_1^2 - c_1), (u_1^3 - 3c_1u_1), (3\lambda(X \circ (u_1^2 - c_1)) - c_2), (u_2u_1), (u_2^2u_1)$$

The Good, the Ugly and the Bad

Concrete example. Let B be a d -dimensional Brownian motion (or a regularisation B^ε) and φ a smooth function. Then $B \in \mathcal{C}^\gamma$ for $\gamma < 1/2$.

$$\varphi(B)DB = \underbrace{\varphi(B) \prec DB}_{\text{the Bad}} + \underbrace{\varphi(B) \circ DB}_{\text{the Ugly}} + \underbrace{\varphi(B) \succ DB}_{\text{the Good, } \mathcal{C}^{2\gamma-1}}$$

and recall the parolinearization

$$\varphi(B) = \varphi'(B) \prec B + \mathcal{C}^{2\gamma}$$

Then

$$\begin{aligned}\varphi(B) \circ DB &= (\varphi'(B) \prec B) \circ DB + \underbrace{\mathcal{C}^{2\gamma} \circ DB}_{\text{OK}} \\ &= \varphi'(B)(B \circ DB) + \mathcal{C}^{3\gamma-1}\end{aligned}$$

Finally

$$\varphi(B)DB = \varphi(B) \prec DB + \varphi'(B) \underbrace{(B \circ DB)}_{\text{"Besov area"}} + \varphi(B) \succ DB + \mathcal{C}^{3\gamma-1}$$

The Besov area

If $d = 1$ (or by symmetrization) we can perform an integration by parts to get

$$B \circ DB = \frac{1}{2}((B \circ DB) + (DB \circ B)) = \frac{1}{2}D(B \circ B)$$

which is well defined and belongs indeed to $\mathcal{C}^{2\gamma-1}$.

In general the Besov area $B \circ DB$ can be defined and studied efficiently using Gaussian arguments:

$$B^\varepsilon \circ DB^\varepsilon \rightarrow B \circ DB$$

almost surely in $\mathcal{C}_{\text{loc}}^{2\gamma-1}$ as $\varepsilon \rightarrow 0$.

Tools: Besov embeddings $L^p(\Omega; C^\theta) \rightarrow L^p(\Omega; B_{p,p}^{\theta'}) \simeq B_{p,p}^{\theta'}(L^p(\Omega))$, Gaussian hypercontractivity $L^p(\Omega) \rightarrow L^2(\Omega)$, explicit L^2 computations.